



Remote conformational responses to enantiomeric excess in carboxylate-binding dynamic foldamers

DOI:

[10.1039/C9CC03895G](https://doi.org/10.1039/C9CC03895G)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Eccles, N., Le Bailly, B., Della Sala, F., Vitorica-yrzebal, I., Clayden, J., & Webb, S. (2019). Remote conformational responses to enantiomeric excess in carboxylate-binding dynamic foldamers. *Chemical Communications*. <https://doi.org/10.1039/C9CC03895G>

Published in:

Chemical Communications

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



COMMUNICATION

Remote conformational responses to enantiomeric excess in carboxylate-binding dynamic foldamers

Received 00th January 20xx,
Accepted 00th January 20xx

Natasha Eccles,^{a,b} Bryden A. F. Le Bailly,^{a,c} Flavio della Sala,^{a,b} Iñigo J. Vitórica-Yrezábal,^a Jonathan Clayden^{c,*} and Simon J. Webb^{a,b,*}

DOI: 10.1039/x0xx00000x

A crystallographically characterised zinc(II)-capped foldamer can sense the enantiomeric excess of scalemic carboxylate solutions, including those produced by enantioselective organocatalysis, and can relay this input signal along the foldamer body to a remote glycinamide group, which then provides an NMR spectroscopic output.

The determination of the enantiomeric excess (*e.e.*) of scalemic product mixtures produced by enantioselective catalysis is a major research field in organic chemistry and the pharmaceutical sciences.^{1,2} Chromatographic separation using chiral solid supports is commonly used to determine *e.e.*, but methods that provide a spectroscopic report of the *e.e.* have the benefits of being faster and cheaper.³

As part of our work using 3_{10} helical α -aminoisobutyric acid oligomers (Aib foldamers) to sense and transmit signals in organic solvents and phospholipid bilayers,⁴ we have become interested in *e.e.* as a signal to be transmitted, particularly as this signal can be generated by an enantioselective catalyst. Previously we showed that capping an Aib octamer with a Cu(II) bis(2-quinolylmethyl)(2-pyridylmethyl)amine (Cu(BQPA)) group gives a synthetic receptor that responds to the binding of a chiral carboxylate ligand (the input signal) by undergoing a conformational change that propagates over several nanometres either in solution or in a lipid bilayer.^{4c} Screening of a series of enantiopure carboxylates showed that the strength of the relayed response depends on the structure of the incoming messenger.^{4c}

Simple Cu(BQPA)²⁺ complexes are able to sense the *e.e.* of scalemic mixtures of chiral carboxylic acids.⁵ The Cu(BQPA)²⁺ complex adopts a helical conformation and coordination to a

chiral carboxylate can favour one conformation over the other. This produces a circular dichroism (CD) signal that, after analysis, indicates the configuration and enantiomeric purity of different acids.⁵ Similarly, zinc(II) complexes of tri(2-pyridylmethyl)amine derivatives have been used to measure the *e.e.* of amino acid or alcohol mixtures.⁶




Figure 1: Structure of Zn(1)²⁺ binding a chiral carboxylate (red, BocPro and BocPip shown below), with the equilibrium between screw senses indicated. Quinoyl groups (blue) and diastereotopic protons (green) on the GlyNH₂ are highlighted.

Although an N-terminal M(BQPA)²⁺ on an Aib foldamer should be able to report on the *e.e.* of a ligand mixture, detection of an *e.e.* “signal” that has been relayed along the length of the foldamer requires a reporter with an orthogonal spectroscopic output. Combining a C-terminal glycinamide (GlyNH₂) with the N-terminal BQPA would provide a receptor with two orthogonal outputs, namely ¹H NMR and CD spectroscopies (Figure 1). The GlyNH₂ group is a particularly well-characterised ¹H NMR reporter group.^{4a,7} The 3_{10} helical conformation of the

^a School of Chemistry, University of Manchester, Oxford Road, Manchester M13 9PL, United Kingdom.

^b Manchester Institute of Biotechnology, University of Manchester, 131 Princess St, Manchester M1 7DN, United Kingdom. E-mail: S.Webb@manchester.ac.uk; Tel: +44 161 306 4524.

^c School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, UK. E-mail: j.clayden@bristol.ac.uk

Electronic Supplementary Information (ESI) available: Synthetic procedures and NMR spectra for new compounds; analytical procedures and data for NMR studies; x-ray crystallography data. See DOI: 10.1039/x0xx00000x

foldamer renders the GlyNH₂ methylene protons diastereotopic, giving an AB system where the chemical shift separation ($\Delta\delta$) is proportional to the excess of one screw sense over the other (the helical excess, *h.e.*);^{7c,8} an achiral foldamer in fast exchange gives apparently isochronous resonances. Unlike CD, this GlyNH₂ reporter is insensitive to absolute screw-sense preference (the major conformer is not identified). Replacement of paramagnetic Cu(II) with diamagnetic Zn(II), combined with use of a short tetrameric Aib foldamer that is in fast *M/P* helical exchange on the NMR spectroscopy timescale,^{8,9} provides the final features found in the receptor Zn(**1**)-2ClO₄ (Figure 1). This synthetic receptor should be able to respond to an input signal, the *e.e.* of a scalemic mixture of carboxylates that could in turn be generated by a catalytic process, and be able to transmit this input signal over the 1 nm length of the foldamer.

BQPA-capped foldamer **1** was synthesised^{4c} and complexed to zinc(II) perchlorate to give Zn(**1**)-2ClO₄. The X-ray crystal structure was determined for Zn(**1**)-2ClO₄ after crystallisation from methanol/diethyl ether (Figure 2). This structure reveals that, as expected, the Aib tetramer body has folded into a short 3_{10} helix. Some Zn(II) complexes of tris(2-pyridylmethyl)amine have been reported to be *pseudo*-octahedrally coordinated¹⁰ with bridging or non-bridging monovalent ligands,¹¹ but the structure of Zn(**1**)-2ClO₄ shows a single water molecule (rather than methanol) coordinated to the Zn(II) centre. This water molecule hydrogen bonds to one of the perchlorate counter ions and then onto the NH of the second Aib residue from the N-terminus. Another network of hydrogen-bonded water molecules also links the bound water molecule to the carbonyl of the BQPA and onto the NH of the third Aib residue. These interactions indicate how the 3_{10} helix might be influenced by chiral ligands bound to the metal centre. Both screw senses are present in the unit cell, with an *M* propeller-shaped conformation in the BQPA associated with a left-handed (*M*) helical sense in the Aib foldamer chain.

Studies of conformational change in analogous Cu(II)(BQPA)-Aib foldamer complexes had shown that amino acid derivatives bearing *N*-alkoxycarbonyl (e.g. Boc protection) or *N*-acyl groups were the most effective.^{4c} In particular, BocPro is a powerful inducer of conformational chirality in the BQPA derivative of an analogous Cu(**1**) foldamer.

Following previous protocols,^{4c} studies of BocPro complexation were performed. The ¹H NMR spectrum after the addition of half an equivalent of Boc-L-Pro to Zn(**1**)-2ClO₄ in CD₃CN (0.015 M, with 1.2 eq. 2,6-lutidine) shows well-resolved and sharp resonances corresponding to Zn(**1**)-2ClO₄ both with and without bound carboxylate (see the ESI, Figure S5), consistent with slow exchange between free and complexed foldamer. The addition of a further 0.5 eq. of Boc-L-Pro led to quantitative formation of the corresponding complex. The resonance from the glycinamide CH₂ shifts upfield by *ca.* 0.25 ppm and splits ($\Delta\delta = 179 \pm 9$ ppb), consistent with an excess of one screw sense over the other.¹² Increasing the concentration of Boc-L-Pro (up to 4 eq.) gave no further significant spectroscopic changes, consistent with 1:1 binding to Boc-L-Pro, and the $\Delta\delta$ of the GlyNH₂ remained constant. The CD

spectrum of the Zn(**1**)(Boc-L-Pro)⁺ complex showed a positive couplet at 238 nm, and fitting of the data from a CD titration to a 1:1 binding model¹³ gave an apparent binding constant of approx. 4×10^6 M⁻¹ (see ESI).




Figure 2: (a,b) X-ray crystal structure of Zn(**1**)-2ClO₄. (a) Plan view showing 3_{10} helix hydrogen bonded to a water bound to the zinc ion. (b) View down the axis of the 3_{10} helix. Selected hydrogen bonds shown to illustrate the hydrogen bonded network from the bound water. The perchlorate furthest from the metal centre is not shown for clarity. Methyl, methylene and methine protons are also not shown for clarity.

Given that complexed and free Zn(**1**)²⁺ were in slow exchange with each other, it was not clear whether the NMR output from the GlyNH₂ reporter would depend on the *e.e.* of the carboxylate ligand mixture. It seemed possible that anisochronous resonances may result even from addition of racemic BocPro: each foldamer complexes a single chiral carboxylate of either D or L configuration, which induces splitting of the same magnitude. However, titration of Zn(**1**) with *rac*-BocPro gave a glycinamide CH₂ resonance that appeared similar to that of Zn(**1**)²⁺ alone, *i.e.* split by the adjacent NH but with no apparent anisochronicity. This suggests that the enantiomeric complexes Zn(**1**)(Boc-L-Pro)⁺ and Zn(**1**)(Boc-D-Pro)⁺ must be exchanging rapidly on the ¹H NMR timescale. Subsequent titration of Zn(**1**)-2ClO₄ (0.015 M in CD₃CN) with scalemic mixtures of BocPro (seven mixtures, 40 to 100% *e.e.*) confirmed this observation and showed that the chemical shift separation ($\Delta\delta$) between the anisochronous signals from the glycinamide methylene of the foldamer was proportional to the *e.e.* of the BocPro. This change in $\Delta\delta$ with *e.e.* mirrors the proportional dependence of ellipticity on *e.e.* (Figure 3), confirming Zn(**1**)²⁺ is a dual CD and NMR sensor.

To explain the dependence of $\Delta\delta$ on *e.e.*, we propose that instead of ligand dissociation and re-association, which is demonstrably slow on the NMR timescale (see ESI and Figure 1), another pathway must exist to facilitate rapid interconversion between coordinated carboxylates. We suggest that fast but weak association of another carboxylate to the zinc(II) centre creates a *pseudo*-octahedral intermediate, which is followed by fast dissociation of one of the complexed carboxylates (see ESI). This proposed intermediate must be at low concentration, since the ¹H NMR spectrum showed no signals that could be ascribed to this complex. Lowering the temperature to -40 °C (see VT-NMR in ESI) provides no more information on the exchange process.

This correlation of *e.e.* and $\Delta\delta$ at the GlyNH₂ terminus provides the opportunity to create a remote reporter of catalytic activity. Zn(**1**)²⁺ can report a signal, namely the *e.e.* of a chiral

carboxylate mixture, much like sensors that use a CD output to report the *e.e.* of a scalemic mixture. However, $\text{Zn}(\mathbf{1})^{2+}$ can relay the signal to a remote site. This *e.e.* signal could be generated by an organocatalytic enantioselective reaction, which would reflect the enantioselectivity of the catalyst.




Figure 3: (a) Partial ^1H NMR spectra (400 MHz, 298 K) showing glycineamide CH_2 region of $\text{Zn}(\mathbf{1})\cdot 2\text{ClO}_4$ in CD_3CN (7.5 μmol in 0.5 mL) mixed with scalemic Boc-Pro (75 mM stock with 1.2 eq. 2,6-lutidine, *e.e.* 0 to 100%). (b) CD spectra of $\text{Zn}(\mathbf{1})\cdot 2\text{ClO}_4$ (0.25 mM) with scalemic BocPro (2 eq.) in MeCN with 2,6-lutidine (2.4 eq.). (c) Change in corrected $\Delta\delta$ (gray) and ellipticity (at $\lambda_{\text{max}} = 238$ nm, black) upon addition of scalemic BocPro (*e.e.* %) to $\text{Zn}(\mathbf{1})\cdot 2\text{ClO}_4$. $\Delta\delta$ values calculated using published methods,^{4a} and were unresolved for *e.e.* < 40%. Linear curve fits shown for ellipticity ($R = 0.99825$) and $\Delta\delta$ ($R = 0.99954$).

The organocatalytic desymmetrisation of anhydrides was selected as a suitable reaction to demonstrate the recognition and relay of enantioselectivity. Chiral amine **4** catalyses the addition of alcohols to cyclic *meso*-anhydrides to give chiral monoesters with high *e.e.*¹⁴ Anhydride **2a** had been desymmetrised in good *e.e.* (85%, predominately the $2R,3S$ isomer) and high yield (90%) by a thiourea analogue of **4**,¹⁵ so desymmetrisation of the same substrate was explored using **4**. Screening indicated that THF gave the best enantioselectivity of those solvents tested (MeOH, 17% *e.e.*; MeCN, 64% *e.e.*). Desymmetrisation of **2a** in THF with methanol (10 eq.) and a catalytic amount of **4** (0.1 eq.) afforded monoester **3a** in good yield (98%) and high enantioselectivity (96% *e.e.*) after 24 h. The same conditions were then applied to anhydride **2b**, albeit with a doubling of the reaction time, and resulted in a similarly high *e.e.* and yield (Figure 4).

The resulting scalemic mixture of **3b**, which has a closer structural analogy with BocPro than **3a**, was then added to a solution of $\text{Zn}(\mathbf{1})^{2+}$ in acetonitrile and the resulting mixture was analysed by CD and NMR spectroscopy. The addition of aliquots of **3b** (1*R*, 2*S*) to $\text{Zn}(\mathbf{1})$ (0.025 mM) gave a positive exciton-coupled circular dichroism (ECCD) signal⁵ at 238 nm in the CD spectra due to an exciton couplet between the

quinoline chromophores. The sign of the change suggests **3b** (1*S*,2*R*), much like (*S*) amino acids, causes a left-handed (*M*) propeller twist in the BQPA.⁵ The intensity of the signal with respect to concentration of **3b** reached a maximum of +19 mdeg after 0.6 eq., perhaps due to the formation of bridged 2:1 foldamer:carboxylate complexes, before declining.




Figure 4: Organocatalytic desymmetrisation of cyclic *meso*-anhydrides by **4** to give scalemic mixtures of carboxylic acids. Enantiomeric excesses (*e.e.*) determined by HPLC on chiral stationary phase (**); conjugation to (*R*)-naphthylethylamine followed by HPLC on chiral stationary phase (*); conjugation to (*R*)-naphthylethylamine followed by ^1H NMR spectroscopy (*). Stereochemistry confirmed (where indicated) by comparison of $[\alpha]_D$ values to reported values. Major isomer shown. # Not separated from anhydride.

For ^1H NMR spectroscopy studies, the monoester **3b** was separated from **4** by washing with 1 M HCl; although **4** did not itself induce splitting in the glycineamide CH_2 resonance, it did obscure key resonances. The unresolved monoester mixture was then dissolved in CD_3CN (75 mM, with 1.2 eq. 2,6-lutidine). Monoester **3b** was added into $\text{Zn}(\mathbf{1})^{2+}$ in CD_3CN (15 mM) to give a 1:2 foldamer:carboxylate ratio and the corresponding ^1H NMR spectrum recorded. A low $\Delta\delta$ value (below 40 ppb) was observed (Table 1). Similarly, **3a** and the cyclohexane analogue of **3b** (**3d**, >98% *e.e.*) both also gave low $\Delta\delta$ values (44 ppb and 39 ppb, see the ESI), and all are much lower than those for Boc-L-Ala, Boc-L-Pro and Boc-L-Pip ($\Delta\delta = 127, 179$ and 231 ppb respectively, Table 1). The racemic mixture from the uncatalysed reaction of **3b** gave a $\Delta\delta$ of 0 ppb, which shows that, as expected, the C-terminal GlyNH₂ reports the *e.e.* of the scalemic mixture, rather than an individual bound signal molecule. The $\Delta\delta$ values do not correlate with the strength of the CD signal, indicating that these foldamers show different analyte selectivity compared to BQPA-based sensors using only a CD output. Since large conformational changes in the BQPA do not necessarily lead to good signal transmission down the helix, this suggests that screw-sense preference may originate from the ligand directly hydrogen-bonding into the Aib backbone.^{4c}

Based upon this hypothesis, a new ligand, **3c**, with a good hydrogen bond acceptor adjacent to the carboxylate was designed. Ligand **3c** is structurally similar to BocPip (Figure 1), which induces a high *h.e.* in Zn(**1**)²⁺ (Table 1). Anhydride **2c** was synthesised from pyridine-2,6-dicarboxylic acid (see the ESI) and subjected to methanolysis catalysed by **4**. Although this substrate was less effectively desymmetrised under standard conditions (25 °C, 1-2 days), a significant increase in the loading of **4** (from 0.1 to 1.1 eq.) and a decrease in reaction temperature (to -20 °C) provides monoester **3c** in good yield and 96% *e.e.*¹⁶ As hoped, **3c** displayed better control over helical sense, although this control ($\Delta\delta = 90$ ppb) was still less than that induced by BocPip ($\Delta\delta = 231$ ppb).¹⁷

Table 1: Table of carboxylate *e.e.* with corresponding ellipticity induced at the N-terminal BQPA (ΔA , ECCD intensity at the first Cotton effect $\lambda_{\max} = 238$ nm) and anisochronicity ($\Delta\delta$) induced in the C-terminal GlyNH₂ of Zn(**1**)-2ClO₄.

Input	<i>e.e.</i>	ΔA^a / mdeg	$\Delta\delta^a$ / ppb
Boc-L-Ala	$\geq 96\%^b$	+43	127 ± 5^c
Boc-L-Pro	$\geq 96\%^b$	+80	179 ± 9^d
Boc-L-Pip	$\geq 97.5\%^b$	+53	231 ± 5^c
3b	$0 \pm 1\%^e$	0	0 ± 5^c
3c	$95 \pm 1\%^e$	+13	29 ± 5^c

^a CD and $\Delta\delta$ values were determined at 25 °C in CD₃CN, 2 eq. of carboxylate. Uncertainties in ΔA are approx. $\pm 10\%$. ^b From suppliers' specifications. ^c Uncertainties in $\Delta\delta$ estimated from NMR line width. ^d From six independent measurements. ^e Uncertainties in *e.e.* estimated from NMR integration.

In summary, a novel Zn(BQPA)-capped dynamic Aib foldamer has been synthesised, structurally characterised and shown to act as a CD and remote NMR sensor of *e.e.* in scalemic mixtures of carboxylates. This foldamer transmits an input signal, the *e.e.* of each scalemic mixture, along the length of foldamer to the reporter group. This type of signal transduction differs from most of those previously reported for Aib foldamers, in that the foldamer senses and responds to a system (an ensemble of signalling molecules)¹⁸ rather than the presence of a specific signalling molecule. By using this signalling relay to sense the outcome from enantioselective organocatalysis, proof-of-concept for signal conversion and amplification has been demonstrated; an unreported chiral signal (catalyst **4**) has been converted into a detectable signal that is amplified and transmitted down the foldamer length.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the BBSRC (DTP studentship), the ERC (Advanced Grant ROCOCO), and the EPSRC (grants EP/P027067/1, EP/N009134/1 and EP/K039547). We thank Mrs R. Sung for HPLC advice. We also thank Dr R. Spiess and the Mass Spectrometry Service for mass spectrometry support.

Notes and references

- (a) D. J. Pollard and J. M. Woodley, *Trends Biotechnol.* 2006, **25**, 66-73. (b) F. G. Mutti, T. Knaus, N. S. Scrutton, M. Breuer and N. J. Turner, *Science*, 2015, **349**, 1525-1529.
- M. J. Gaunt, C. C. C. Johansson, A. McNally and N. T. Vo, *Drug Discov. Today*, 2007, **12**, 8-27.
- (a) D. Leung, S. O. Kang and E. V. Anslyn, *Chem. Soc. Rev.*, 2012, **41**, 448-479. (b) A. Akdeniz, T. Minami, S. Watanabe, M. Yokoyama, T. Ema and P. Anzenbacher, Jr, *Chem. Sci.*, 2016, **7**, 2016-2022.
- (a) R. A. Brown, V. Diemer, S. J. Webb and J. Clayden, *Nature Chem.*, 2013, **5**, 853-860. (b) M. De Poli, W. Zawodny, O. Quinonero, M. Lorch, S. J. Webb and J. Clayden, *Science* 2016, **352**, 575-580. (c) F. G. A. Lister, B. A. F. Le Bailly, S. J. Webb and J. Clayden, *Nature Chem.* 2017, **9**, 420-425. (d) F. G. A. Lister, N. Eccles, S. J. Pike, R. A. Brown, G. F. S. Whitehead, J. Raftery, S. J. Webb and J. Clayden, *Chem. Sci.* 2018, **9**, 6860-6870.
- L. A. Joyce, M. S. Maynor, J. M. Dagna, G. M. da Cruz, V. M. Lynch, J. W. Canary and E. V. Anslyn, *J. Am. Chem. Soc.*, 2011, **133**, 13746-13752.
- (a) F. A. Scaramuzza, G. Licini, and C. Zonta, *Chem. Eur. J.* 2013, **19**, 16809-16813. (b) L. You, G. Pescitelli, E. V. Anslyn and L. Di Bari, *J. Am. Chem. Soc.* 2012, **134**, 7117-7125.
- (a) R. A. Brown, T. Marcelli, M. De Poli, J. Solà and J. Clayden, *Angew. Chem. Int. Ed.*, 2012, **51**, 1395-1399. (b) M. De Poli, L. Byrne, R. A. Brown, J. Solà, A. Castellanos, T. Boddaert, R. Wechsel, J. D. Beadle and J. Clayden, *J. Org. Chem.*, 2014, **79**, 4659-4675. (c) D. Mazzier, M. Crisma, M. De Poli, G. Marafon, C. Peggion, J. Clayden and A. Moretto, *J. Am. Chem. Soc.*, 2016, **138**, 8007-8018. (d) S. J. Pike, M. De Poli, W. Zawodny, J. Raftery, S. J. Webb and J. Clayden, *Org. Biomol. Chem.*, 2013, **11**, 3168-3176.
- J. Solà, G. A. Morris and J. Clayden, *J. Am. Chem. Soc.* 2011, **133**, 3712-3715.
- R.-P. Hummel, C. Toniolo and G. Jung, *Angew. Chem. Int. Ed.* 1987, **26**, 1150-1152.
- (a) E. Asato, S. Miyazato, H. Tohma, S. Takara, M. Tadokoro, Y. Miyazato, D. Yoshioka and M. Mikuriya, *Chem. Lett.*, 2009, **38**, 1170-1171. (b) L. Metteau, S. Parsons and J. C. Mareque-Rivas, *Inorg. Chem.*, 2006, **45**, 6601-6603. (c) A. Wada, S. Yamaguchi, K. Jitsukawa and H. Masuda, *Angew. Chem. Int. Ed.*, 2005, **44**, 5698-5701. (d) L. Xue, H.-H. Wang, X.-J. Wang and H. Jiang, *Inorg. Chem.*, 2008, **47**, 4310-4318.
- (a) N. N. Murthy and K. D. Karlin, *Chem. Commun.*, 1993, 1236-1238. (b) C. Duboc, T. Phoeung, D. Jouvenot, A. G. Blackman, L. F. McClintock, J. Pecaut, A. Deronzier and M.-N. Collomb, *Polyhedron*, 2007, **26**, 5243-5249. (c) Y. Mikata, K. Kawata, S. Takeuchi, K. Nakanishi, H. Konno, S. Itami, K. Yasuda, S. Tamotsu and S. C. Burdette, *Dalton Trans.*, 2014, **43**, 10751-10759.
- The use of a ¹³C reporter at the C-terminus shows that Boc-L-Pro gives a *P* helical preference. See B. A. F. Le Bailly, PhD thesis, University of Manchester, 2015.
- P. Kuzmic, *Anal. Biochem.*, 1996, **237**, 260-273.
- S. H. Oh, H. S. Rho, J. W. Lee, J. E. Lee, S. H. Youk, J. Chin and C. E. Song, *Angew. Chem. Int. Ed.* 2008, **47**, 7872-7875.
- A. Peschiulli, Y. Gun'ko and S. J. Connon, *J. Org. Chem.*, 2008, **73**, 2454-2457.
- The configuration of the major enantiomer is unknown.
- Residual catalyst in the **3c** mixture interfered with the measurement of the BQPA ECCD signal from Zn(**1**)-2ClO₄.
- J. Brioché, S. J. Pike, S. Tshpelevitsh, I. Leito, G. A. Morris, S. J. Webb and J. Clayden, *J. Am. Chem. Soc.* 2015, **137**, 6680-6691.