



# An observational cohort study investigating risk of malnutrition using the Malnutrition Universal Screening Tool in patients with stroke.

DOI:

[10.1016/j.jstrokecerebrovasdis.2019.104405](https://doi.org/10.1016/j.jstrokecerebrovasdis.2019.104405)

## Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

## Citation for published version (APA):

Sremanakova, J., Burden, S., Kama, Y., Gittins, M., Lal, S., Smith, C., & Hamdy, S. (2019). An observational cohort study investigating risk of malnutrition using the Malnutrition Universal Screening Tool in patients with stroke. *Journal of Stroke and Cerebrovascular Diseases*.  
<https://doi.org/10.1016/j.jstrokecerebrovasdis.2019.104405>

## Published in:

Journal of Stroke and Cerebrovascular Diseases

## Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

## General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

## Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact [uml.scholarlycommunications@manchester.ac.uk](mailto:uml.scholarlycommunications@manchester.ac.uk) providing relevant details, so we can investigate your claim.



1 **ABSTRACT**

2 **Background:** Malnutrition in patients hospitalised with a stroke have been assessed using  
3 different nutritional screening methods but there is a paucity of data linking risk of  
4 malnutrition to clinical outcomes using a validated tool.

5 **Aims:** To identify the prevalence of malnutrition risk in patients after a stroke and assess the  
6 predictive value of the Malnutrition Universal Screening Tool on clinical outcomes.

7 **Patients and methods:** Using data from electronic records and the Sentinel Stroke National  
8 Audit Programme (January 2013 and March 2016), patients aged > 18 years with confirmed  
9 stroke admitted to a tertiary care stroke unit were assessed for risk of malnutrition. The  
10 association between malnutrition risk and clinical outcomes was investigated and adjusted  
11 for confounding variables.  
12

13 **Results:** Of 1101 patients, 66% were screened at admission. Most patients (n= 571, 78.5%)  
14 were identified as being at low risk, 4.1% (n=30) at medium risk and 17.4% (n=126) at high  
15 risk of malnutrition. Compared with low risk, patients with medium or high risk of  
16 malnutrition were more likely to have a longer hospital stay (IRR 1.30, 95% CI 1.07, 1.58),  
17 and had greater risk of mortality (10.9% versus 3.5%, 95% CI 0.03, 0.13).

18 **Conclusions:** Prevalence of malnutrition assessed by Malnutrition Universal Screening Tool  
19 in patients after a stroke was relatively low, but nearly a third of patients were not  
20 screened. Patients classified as being at medium or high risk of malnutrition were more  
21 likely to experience negative outcomes. Early identification of this population may improve  
22 outcome if appropriate care is provided.

23  
24  
25  
26  
27  
28

29 **Introduction**

30 Early post-stroke complications include dysphagia, infections, recurrence of stroke, increased  
31 intracranial pressure and malnutrition. Documented prevalence of malnutrition in patients  
32 after a stroke ranges from 6.1% to 62% (1) and has been shown to affect clinical outcomes (2,  
33 3). However, in the majority of patients, malnutrition is considered an avoidable consequence  
34 of a stroke (3). The most obvious indication of malnutrition is rapid weight loss and this  
35 pattern of weight change is recognised as an important predictor of poor outcomes (4).  
36 Available evidence indicates that, after a stroke, a reduction in weight of more than 3 kg both  
37 in the short and long term can impact on risk of mortality (5). Importantly, studies indicate  
38 that nutritional interventions can help improve patients' clinical outcomes if initiated early  
39 (6-8).

40

41 To provide nutritional interventions to patients at risk of malnutrition, those who are  
42 malnourished need to be identified. This can be achieved with the use of a nutritional  
43 screening tool, as recommended by national clinical guidelines for stroke which advise  
44 screening for malnutrition risk on admission and at least weekly thereafter by trained staff  
45 using a structured tool (9). A number of studies have reported on nutritional status in the  
46 stroke population using different assessment methods or screening tools (2, 6, 8, 10-15).  
47 However, a large discrepancy exists in the tools used, time of assessment and definition of  
48 malnutrition.

49

50 There is a difference between nutritional screening and nutritional assessment (16).  
51 Nutritional screening tools are designed to be used as rapid and simple methods to identify  
52 those at risk of malnutrition, while nutritional assessment involves comprehensive  
53 assessment of nutritional status by trained personnel to identify malnutrition, which can then  
54 be treated with appropriate nutritional interventions (17). The most frequently used  
55 screening tool in the UK is the Malnutrition Universal Screening Tool (MUST), a feasible way  
56 of identifying patients at risk of malnutrition in clinical and community settings (18), which  
57 has been included in the Global Leadership Initiative on Malnutrition (GLIM), a new consensus  
58 on assessment of malnutrition (19). Although MUST has been validated against nutritional  
59 assessment methods and other screening tools (20) in mixed patient populations, the  
60 evidence of its predictive value in patients after a stroke is thus far unclear.

61 The paucity and inconsistency of evidence on the prevalence of malnutrition in patients after  
62 a stroke, leaves clinical practitioners with uncertainty. Therefore, the aim of this observational  
63 clinical registry and audit was to identify the prevalence of risk of malnutrition in patients  
64 after a stroke determined by MUST in a UK based secondary care facility and its association  
65 with clinical outcomes including mortality and length of stay (LoS).

## 66 **Materials and Methods**

67 This was a retrospective analysis of prospectively collected data from local Sentinel Stroke  
68 National Audit Programme (SSNAP) and the electronic patient record (EPR) at the Greater  
69 Manchester Comprehensive Stroke Centre (CSC) based at Salford Royal Foundation Trust. All  
70 patients with suspected stroke within 48-hours of onset were admitted to the stroke unit  
71 based on location of home residence. Patients with a confirmed stroke, aged at least 18 years  
72 and with a Salford postcode receiving their ongoing care at CSC were eligible for the study,  
73 and identified using Salford SSNAP data. Core SSNAP data were extracted and merged with  
74 additional data from the EPR. The clinical parameters included patients' characteristics,  
75 baseline nutritional status including weight and body mass index (BMI); MUST score at  
76 admission and before discharge; mortality, National Institutes of Health Stroke Scale (NIHSS  
77 scale), Modified Rankin Scale (mRS), complications and LoS. Ethical approval for this  
78 secondary data analysis was sought from the University of Manchester's Ethics Committee  
79 (UREC). As all data were collected as part of routine clinical care, following (proportionate)  
80 review and consideration by UREC, the study was given exemption from the requirement for  
81 ethical approval as a clinical registry, service evaluation and audit. The extraction of data was  
82 performed by the Information Technology Department at Salford Royal NHS Foundation Trust  
83 using the Electronic Patient Records (EPR). These data were matched with that of the Sentinel  
84 Stroke National Audit Programme (SSNAP) database by a member of the clinical team and  
85 fully anonymised. The research team had access to the anonymised data only. The study was  
86 formally registered as a clinical audit at the Clinical Audit Department (reference number  
87 2016151) at Salford Royal NHS Foundation Trust.

88

## 89 **MUST score**

90 MUST score were collected from SSNAP data. Healthcare professionals performed MUST  
91 assessment as part of routine hospital admissions procedure and before discharge. The MUST

92 methodology used incorporates three independent variables: *BMI score* (BMI >20.0= 0, BMI  
93 18.5-20.0= 1, BMI < 18.5= 2); *unplanned weight loss* in previous 3-6 months (weight loss  
94 <5%=0, weight loss 5-10% = 1 and weight loss >10 % = 2); and *acute disease effect score* (a  
95 score of 2 was added if a patient was recently affected by a disease and there was no  
96 nutritional intake or likely to be no nutritional intake for more than 5 days). For BMI and  
97 weight loss, each variable is scored on a scale of 0 to 2. A total sum of scores is used to  
98 categorise the risk of malnutrition as low (0), medium (1) and high ( $\geq 2$ ) (21). Details on MUST  
99 score assessment can be found at the British Association for Parenteral and Enteral  
100 Nutrition website (22). We further categorised MUST into a dichotomised variable (low risk  
101 of malnutrition 0, versus medium to high risk of malnutrition  $\geq 1$ ).

102

### 103 **Statistical methods**

104 Standard descriptive statistics summarised the patient and clinical characteristics. Statistical  
105 inference determined the association between prevalence of risk for malnutrition and clinical  
106 outcomes, whilst adjusting for potential confounders in a multiple logistic regression analysis.  
107 LoS defined as a count of the days as inpatient was modelled using a negative binomial  
108 regression to investigate the relationship between LoS and MUST. Confounders were  
109 identified prior to analysis. Weight, BMI and mRS were not included due to collinearity. The  
110 analysis was adjusted for age, gender, type of stroke, NIHSS scale and comorbidities. Further  
111 (logistic) regression modelling investigating the association between MUST and deaths in  
112 hospital, however deaths at 6-months from admission was not possible due to limited death  
113 events occurring in those with a MUST score. A Fishers exact test however was possible for  
114 death at 6 months and is reported.

115

### 116 **Results**

117 Between January 2013 and March 2016, records of 1,101 patients who met the eligibility  
118 criteria were extracted from EPR and combined with corresponding SSNAP data. The patient's  
119 baseline characteristics based on dichotomised MUST score for the total population are  
120 shown in Table 1, and mRS at pre-admission and discharge by MUST category is presented at  
121 Figure 1. Mean age was 73.6 (SD 13.6) years and 94% of patients included were of a Caucasian  
122 origin with equal gender distribution. Just over half of patients (n=563, 51.2%) had a history

123 of hypertension; about a fifth of patients (n=209, 19%) had diabetes mellitus, atrial fibrillation  
124 (n=197, 17.9%), and a small proportion had (n=51, 4.4%) congestive heart failure. The vast  
125 majority of patients (n=976, 89.2%) were diagnosed with ischemic stroke, with the remainder  
126 (n=118, 10.8%) being diagnosed with a primary intracerebral haemorrhage. A fifth of patients  
127 had a previous transient ischemic attack (n=218, 19.8%). Based on NIHSS scale(23), no stroke  
128 symptoms (score 0) were present in 10.8% of patients, minor stroke (score 1 to 4) were  
129 experienced by 40.2%, moderate stroke (score 5-15) by 33.5%, moderate to severe stroke  
130 (score 16-20) by 7.3% and severe stroke (score 21 to 42) by 8.3% patients (Table 1).

131

132 **Table 1**

133 **Figure 1**

134

135 *MUST score*

136 A MUST score was recorded in 66% of patients both on admission and prior to discharge from  
137 hospital. On admission, the majority of patients 78.5% (n=571) were at low risk of  
138 malnutrition (MUST score 0), 4.1% (n=30) were considered to be at medium risk (MUST score  
139 1) and 17.4% (n=126) were classified as being at high risk of malnutrition (MUST score  $\geq 2$ ).  
140 At the point of discharge from hospital, the number of patients with low risk of malnutrition  
141 increased to 85.2% (n=618), the number of patients with medium risk increased to 5.8 %  
142 (n=42) and the number of patients in high risk group decreased to 9% (n=65). As expected, 38  
143 (90.5%) of patients with BMI between 18.5-20 kg/m<sup>2</sup> and 18 (100%) of patients with BMI less  
144 than 18 kg/m<sup>2</sup> were identified as being at medium to high risk of malnutrition .

145

146 *Negative binomial regression*

147 Table 2 reports the incidence rate ratios (IRR) and corresponding 95% confidence intervals for  
148 the adjusted association between LoS and risk of malnutrition defined by a dichotomised  
149 MUST score. Compared to low risk, medium or high risk of malnutrition was associated with  
150 an IRR = 1.30 (95% CI 1.07 to 1.58) indicating a 30% increase in the risk of a longer hospital  
151 stay. Additionally, a linear increase in risk of a longer hospital stay was observed for NIHSS  
152 scores increasing in severity from 'moderate' through to 'severe' (IRR = 1.49, 2.64 and 3.14)  
153 when compared to no symptoms. Interestingly, minor stroke was associated with a 30%

154 decrease compared to no symptoms (IRR = 0.72, 95% CI 0.56 to 0.93). Of the co-morbidities,  
155 only diabetes was associated with an increase in LoS of 31% (IRR = 1.31, 95% CI 1.07 to 1.61).

156

## 157 **Table 2**

158

### 159 *LoS and complications*

160 Median LoS for all patients was 7-days and ranged between 0 to 147 days. Twenty patients  
161 (2%) were diagnosed with urinary tract infection in hospital. Pneumonia affected 53 patients  
162 (5.4%). By comparison, patients who were not assessed for MUST (n=374), had a median LoS  
163 of 6-days and ranged between 0 to 147 days. Urinary tract infections were identified in 3  
164 (0.8%) and pneumonia in 33 (8.8%) of the non-MUST assessed patients (Table 3).

165

### 166 *Mortality*

167 From the total sample of 1,101 patients, 214 (19.4%) patients had died at 6-months follow  
168 up, and from these, 161 (14.6%) died in hospital. From 161 patients who died in hospital, only  
169 one patient had been assessed for malnutrition with MUST score on admission. Survival of  
170 patients who died in hospital was median 3.5 (range 0-147) days. Of these, 68 (42.5%) patients  
171 did not survive for longer than a day. For patients with a MUST  $\geq 1$  (indicating a medium to  
172 high risk of malnutrition), the number of deaths was greater compared to those with a MUST=  
173 0 (indicating a low risk of malnutrition), 17/159 (10.9%) versus n= 20/571 (3.5%), respectively,  
174 P=0.001. Main cause of the death at six months was recorded in 149 out of 214 patients, an  
175 ischemic stroke in 44% (n=66), spontaneous cerebral haemorrhage 16% (n=24), pneumonia  
176 in 14% (n=21), cancer 11% (n=16) and other in 15% (n=22) of patients (Table 3).

177

## 178 **Table 3**

### 179 **Discussion**

180 In this observational study we examined the prevalence of risk for malnutrition in patients  
181 after a stroke and described the use of MUST as a previously validated screening tool (20) to  
182 determine future risk of malnutrition. From a total population of 1,101 patients admitted to  
183 the hospital during a three year period, only two-thirds were screened using MUST. A similar  
184 proportion of patients screened for malnutrition was previously reported by the Food Trial

185 Collaboration (6). This indicates that around one third of patients with a potential risk of  
186 malnutrition are not captured by routine screening. Due to the records based dataset in the  
187 form of EPR data merged with SSNAP, we were unable to ascertain whether the reduced  
188 numbers of patients screened is related to lack of assessment or missing data.

189

190 In our sample screened for malnutrition, 21.5% of people admitted to the hospital with  
191 stroke were at moderate to high risk of malnutrition. However, due to the paucity of  
192 evidence on risk of malnutrition assessed by MUST in patients after stroke, our data can  
193 only be compared with one other study of a similar design that showed 36% risk of  
194 malnutrition (24). Surprisingly, we observed a higher proportion of patients who were at  
195 low risk of malnutrition at discharge compare to admission. This decrease (in the high risk  
196 prevalence) might reflect good nutritional care provided to patients prior to hospital  
197 discharge at our centre. A further investigation detailing the care pathway might provide  
198 more insight into observed more favourable rate of malnutrition risk.

199

200 The results of this study concur with previous data demonstrating the ability of MUST to  
201 independently predict negative outcomes in patients after a stroke (24), raising the possibility  
202 that intervening may improve clinical outcomes. Also, it was previously shown that up to one  
203 quarter of patients after a stroke became malnourished within the first week of admission,  
204 and the risk increased with prolonged hospital stay (10, 15). In our study, a median length of  
205 stay of 7-days was lower than 13-days and 25-days previously reported (6, 25). This was  
206 possibly affected by a relatively low prevalence of malnutrition in the population.

207

208 Although the level of malnutrition in patients after admission with stroke is relatively low in  
209 comparison to other disease states (26), possibly reflecting their acute presentation, these  
210 patients still require clinical recognition and support, since malnutrition has been frequently  
211 reported as an independent prognostic factor of post-stroke complications and poor clinical  
212 outcomes (2, 14, 15). The importance of addressing malnutrition is underlined by the cost of  
213 hospitalisation for patients with stroke, which was estimated to be up to 77% higher in  
214 patients who have a high risk of malnutrition compared to those patients who have a low risk  
215 (13, 24). Thus, early identification of patients at risk and provision of appropriate nutritional



216 support might help not only to reduce poorer clinical outcomes in patients but also reduce  
217 the burden on financial resources.

218

219 As with all retrospective studies, the main limitation of this study was the use of data collected  
220 for clinical audit and patient's records. We were limited in records availability and consequent  
221 missing data for individual outcomes. Also, the impact of dysphagia and post-stroke  
222 depression, both linked to oral intake and nutritional status were not directly factored into  
223 our analyses, which may have impacted on outcomes. Data were analysed from a single  
224 hospital in the UK and the rates of screening may reflect a degree of selection bias. Moreover  
225 the MUST score was recorded in only two thirds of patients admitted to hospital with a stroke,  
226 which limits our ability to draw strong conclusions on prevalence of malnutrition and its  
227 association with clinical outcomes. However, despite the limitations, these data provide an  
228 insight into a risk of malnutrition in a large number of participants after a stroke.

229

### 230 **Summary and Conclusion**

231 Our findings show that there is a significant group of patients at risk of malnutrition, and  
232 malnutrition risk as identified by MUST is associated with longer hospital stay even after  
233 adjustment for confounders. Hence, these patients require clinical recognition, and further  
234 prospective studies are warranted to evaluate any effect of nutritional interventions.  
235 Importantly, our data also show that there are substantial numbers of patients who are not  
236 screened for malnutrition on admission to hospital. These patients might be at risk of  
237 malnutrition but remain unidentified.

238

239

240

241

242

243

244

245

246

247

248 **References**

- 249 1. Foley NC, Salter KL, Robertson J, Teasell RW, Woodbury MG. Which reported estimate of the  
250 prevalence of malnutrition after stroke is valid? *Stroke*. 2009;40(3):e66-74.
- 251 2. Martineau J, Bauer JD, Isenring E, Cohen S. Malnutrition determined by the patient-  
252 generated subjective global assessment is associated with poor outcomes in acute stroke patients.  
253 *Clinical nutrition*. 2005;24(6):1073-7.
- 254 3. Bouziana SD, Tziomalos K. Malnutrition in patients with acute stroke. *Journal of nutrition*  
255 *and metabolism*. 2011;2011:167898.
- 256 4. Martin L, Senesse P, Gioulbasanis I, Antoun S, Bozzetti F, Deans C, et al. Diagnostic criteria  
257 for the classification of cancer-associated weight loss. *Journal of clinical oncology : official journal of*  
258 *the American Society of Clinical Oncology*. 2015;33(1):90-9.
- 259 5. Jonsson AC, Lindgren I, Norrving B, Lindgren A. Weight loss after stroke: a population-based  
260 study from the Lund Stroke Register. *Stroke*. 2008;39(3):918-23.
- 261 6. Food Trial Collaboration. Poor nutritional status on admission predicts poor outcomes after  
262 stroke: observational data from the FOOD trial. 2003 Jun. Report No.: 1524-4628 (Electronic)  
263 0039-2499 (Linking) Contract No.: 6.
- 264 7. Axelsson K, Asplund K, Norberg A, Alafuzoff I. Nutritional status in patients with acute stroke.  
265 *Acta medica Scandinavica*. 1988;224(3):217-24.
- 266 8. Gariballa SE, Parker SG, Taub N, Castleden CM. Influence of nutritional status on clinical  
267 outcome after acute stroke. *The American journal of clinical nutrition*. 1998;68(2):275-81.
- 268 9. Rudd A, Bowen A, Young G, James M. National clinical guideline for stroke: 5th edition 2016.  
269 *Clinical Medicine*. 2017 Feb 7:69-71.
- 270 10. Davalos A, Ricart W, Gonzalez-Huix F, Soler S, Marrugat J, Molins A, et al. Effect of  
271 malnutrition after acute stroke on clinical outcome. *Stroke*. 1996;27(6):1028-32.
- 272 11. Davis JP, Wong AA, Schluter PJ, Henderson RD, O'Sullivan JD, Read SJ. Impact of pre-morbid  
273 undernutrition on outcome in stroke patients. *Stroke*. 2004;35(8):1930-4.
- 274 12. Chai J, Chu FC, Chow TW, Shum NC. Prevalence of malnutrition and its risk factors in stroke  
275 patients residing in an infirmary. *Singapore medical journal*. 2008;49(4):290-6.
- 276 13. Lim SL, Ong KC, Chan YH, Loke WC, Ferguson M, Daniels L. Malnutrition and its impact on  
277 cost of hospitalization, length of stay, readmission and 3-year mortality. *Clinical nutrition*.  
278 2012;31(3):345-50.
- 279 14. Shen HC, Chen HF, Peng LN, Lin MH, Chen LK, Liang CK, et al. Impact of nutritional status on  
280 long-term functional outcomes of post-acute stroke patients in Taiwan. *Archives of gerontology and*  
281 *geriatrics*. 2011;53(2):e149-52.
- 282 15. Yoo SH, Kim JS, Kwon SU, Yun SC, Koh JY, Kang DW. Undernutrition as a predictor of poor  
283 clinical outcomes in acute ischemic stroke patients. *Archives of neurology*. 2008;65(1):39-43.
- 284 16. Charney P. Nutrition screening vs nutrition assessment: how do they differ? *Nutrition in*  
285 *clinical practice : official publication of the American Society for Parenteral and Enteral Nutrition*.  
286 2008;23(4):366-72.
- 287 17. Young AM, Kidston S, Banks MD, Mudge AM, Isenring EA. Malnutrition screening tools:  
288 comparison against two validated nutrition assessment methods in older medical inpatients.  
289 *Nutrition*. 2013;29(1):101-6.
- 290 18. Elia M, Russell C. Nutrition screening survey in the UK and Republic of Ireland in 2011  
291 BAPEN. 2012.
- 292 19. Cederholm T, Jensen GL, Correia MITD, Gonzalez MC, Fukushima R, Higashiguchi T, et al.  
293 GLIM criteria for the diagnosis of malnutrition – A consensus report from the global clinical nutrition  
294 community. *Clinical nutrition*. 2018.
- 295 20. Stratton RJ, Hackston A, Longmore D, Dixon R, Price S, Stroud M, et al. Malnutrition in  
296 hospital outpatients and inpatients: prevalence, concurrent validity and ease of use of the  
297 'malnutrition universal screening tool' ('MUST') for adults. *The British journal of nutrition*.  
298 2004;92(5):799-808.

- 299 21. Elia M. THE 'MUST' REPORT Nutritional screening of adults: a multidisciplinary responsibility  
300 Redditch: BAPEN 2003 [
- 301 22. BAPEN. Malnutrition Universal Screening Tool2003 May 2019; (May ). Available from:  
302 [https://www.bapen.org.uk/pdfs/must/must\\_full.pdf](https://www.bapen.org.uk/pdfs/must/must_full.pdf).
- 303 23. Hage V. The NIH stroke scale: a window into neurological status. NurseCom Nursing  
304 Spectrum (Greater Chicago). 2011;24(15):44-9.
- 305 24. Gomes F, Emery PW, Weekes CE. Risk of Malnutrition Is an Independent Predictor of  
306 Mortality, Length of Hospital Stay, and Hospitalization Costs in Stroke Patients. Journal of stroke and  
307 cerebrovascular diseases : the official journal of National Stroke Association. 2016;25(4):799-806.
- 308 25. Ha L, Hauge T, Spenning AB, Iversen PO. Individual, nutritional support prevents  
309 undernutrition, increases muscle strength and improves QoL among elderly at nutritional risk  
310 hospitalized for acute stroke: a randomized, controlled trial. Clinical nutrition. 2010;29(5):567-73.
- 311 26. Russell C, Elia M. Nutrition screening survey in hospitals in the UK, 2007-2011. BAPEN. 2014.  
312