

Table S1. Reported incidences of eclampsia in selected countries within each continent.

		Years	Original reported	Reference
AMERICA	Colombia	1998	8.1 in 1000 deliveries	(Conde-Agudelo and Kafury-Goeta, 1998)
	Colombia, Ecuador, El Salvador, Panama, Peru, and Venezuela	2012	1 in 587 deliveries.	(Vigil-De Gracia et al., 2015)
	USA	1996-2004	8.2 in 10.000 deliveries	(Hutcheon et al., 2011)
	Canada	2007	5.7 in 10.000 deliveries	(Hutcheon et al., 2011)
EUROPE	Spain	2001-2003	1 in 2546 pregnancies	(Cortes-Perez et al., 2009)
	UK	2005	2.7 in 10.000 deliveries	(Hutcheon et al., 2011)
	Finland	2006-2010	1.5 in 10.000 deliveries	(Jaatinen and Ekholm, 2016)
	Denmark, Norway, Sweden	1998-2000	5 in 10.000 deliveries	(Hutcheon et al., 2011)
	Netherlands	1996-2004	6 in 10.000 deliveries	(Hutcheon et al., 2011)
ASIA	Taiwan	1998-2010	32 in 100.000 pregnancies.	(Chan et al., 2015)
	Bangladesh	1998-2000	9 in 100 pregnancies	(Begum et al., 2004)
	Singapore	1999-2003	0.02 in 100 pregnancies	(Tan et al., 2006)
	India	2014	1.49 in 100 deliveries	(Nobis and Hajong, 2016)
AFRICA	Nigeria	2008-2014	1.12 in 100 deliveries	(Esike et al., 2017)
OCEANIA	Australia	2000-2008	8.6 in 10.000 pregnancies	(Thornton et al., 2013)

- Begum, M.R., Begum, A., Quadir, E., Akhter, S., and Shamsuddin, L. (2004). Eclampsia: still a problem in Bangladesh. *MedGenMed* 6(4), 52.
- Chan, T.F., Tung, Y.C., Wang, S.H., Lee, C.H., Lin, C.L., and Lu, P.Y. (2015). Trends in the incidence of pre-eclampsia and eclampsia in Taiwan between 1998 and 2010. *Taiwan J Obstet Gynecol* 54(3), 270-274. doi: 10.1016/j.tjog.2013.06.021.
- Conde-Agudelo, A., and Kafury-Goeta, A.C. (1998). Epidemiology of eclampsia in Colombia. *Int J Gynaecol Obstet* 61(1), 1-8.
- Cortes-Perez, S., Perez Milan, F., Gobernado Tejedor, J., and Mora Cepeda, P. (2009). Epidemiologia de los estados hipertensivos del embarazo. *Clin Invest Gin Obst* 36(4), 132.
- Esike, C.O.U., Chukwuemeka, U.I., Anozie, O.B., Eze, J.N., Aluka, O.C., and Twomey, D.E. (2017). Eclampsia in rural Nigeria: The unmitigating catastrophe. *Ann Afr Med* 16(4), 175-180. doi: 10.4103/aam.aam_46_16.
- Hutcheon, J.A., Lisonkova, S., and Joseph, K.S. (2011). Epidemiology of pre-eclampsia and the other hypertensive disorders of pregnancy. *Best Pract Res Clin Obstet Gynaecol* 25(4), 391-403. doi: 10.1016/j.bpobgyn.2011.01.006.
- Jaatinen, N., and Ekholm, E. (2016). Eclampsia in Finland; 2006 to 2010. *Acta Obstet Gynecol Scand* 95(7), 787-792. doi: 10.1111/aogs.12882.
- Nobis, P.N., and Hajong, A. (2016). Eclampsia in India Through the Decades. *J Obstet Gynaecol India* 66(Suppl 1), 172-176. doi: 10.1007/s13224-015-0807-5.
- Tan, K.H., Kwek, K., and Yeo, G.S. (2006). Epidemiology of pre-eclampsia and eclampsia at the KK Women's and Children's Hospital, Singapore. *Singapore Med J* 47(1), 48-53.
- Thornton, C., Dahlen, H., Korda, A., and Hennessy, A. (2013). The incidence of preeclampsia and eclampsia and associated maternal mortality in Australia from population-linked datasets: 2000-2008. *Am J Obstet Gynecol* 208(6), 476 e471-475. doi: 10.1016/j.ajog.2013.02.042.
- Vigil-De Gracia, P., Rojas-Suarez, J., Ramos, E., Reyes, O., Collantes, J., Quintero, A., et al. (2015). Incidence of eclampsia with HELLP syndrome and associated mortality in Latin America. *Int J Gynaecol Obstet* 129(3), 219-222. doi: 10.1016/j.ijgo.2014.11.024.