



# Minority Status and Mental Distress – a comparison of group density effects

DOI:

[10.1017/S0033291716001835](https://doi.org/10.1017/S0033291716001835)

## Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

## Citation for published version (APA):

Schofield, P., Das-Munshi, J., Becares, L., Morgan, C., Bhavsar, V., Hotopf, M., & Hatch, S. (2016). Minority Status and Mental Distress – a comparison of group density effects. *Psychological Medicine*.  
<https://doi.org/10.1017/S0033291716001835>

## Published in:

Psychological Medicine

## Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

## General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

## Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact [uml.scholarlycommunications@manchester.ac.uk](mailto:uml.scholarlycommunications@manchester.ac.uk) providing relevant details, so we can investigate your claim.



# Minority Status and Mental Distress – a comparison of group density effects

Schofield, P; Das-Munshi, J; Bécares, L; Morgan, C; Bhavsar, V; Hotopf, M; Hatch, SL

## Abstract

### Background

It has been observed that mental disorders, such as psychosis, are more common for people in some ethnic groups in areas where their ethnic group is less common. We set out to test whether this ethnic density effect reflects minority status in general, by looking at three situations where individual characteristics differ from what is usual in a locality.

### Methods

Using data from the South East London Community Health study (n=1698) we investigated associations between minority status (defined by: ethnicity, household status and occupational social class) and risk of psychotic experiences, common mental disorders and parasuicide. We used a multilevel logistic model to examine cross-level-interactions between minority status at individual and neighbourhood levels.

### Results

Being Black in an area where this was (10%) less common was associated with higher odds of psychotic experiences (odds ratio (OR) 1.34 95% CI 1.07 to 1.67), and attempted suicide (OR 1.84 95% CI 1.19 to 2.85). Living alone where this was less usual (10% less) was associated with increased odds of psychotic experiences (OR 2.18 95% CI 0.91 to 5.26), while being in a disadvantaged social class where this was less usual (10% less) was associated with increased odds of attempted suicide (OR 1.33 95% CI 1.03 to 1.71). We found no evidence for an association with common mental disorders.

### Conclusions

The relation between minority status and mental distress was most apparent when defined in terms of broad ethnic group but was also observed for individual household status and occupational social class.

4,784 words

## Introduction

Having a minority status, in terms of a defining social characteristic that differs from others in a locality, has been associated with an increased risk of mental illness (Shaw et al., 2012; van Os et al., 2010; Zammit et al., 2010). Most studies, to date, define minority status in terms of ethnicity where, as the proportion of ethnic minorities in a locality increases – and thus their minority status decreases – the risk of mental illness is reduced. This “ethnic density” effect is well documented (Boydell et al., 2001; Kirkbride et al., 2007; Stafford et al., 2009; Veling et al., 2008). A number of explanations for the ethnic density effect have been proposed including: the absence of social support and social capital, greater vulnerability to discrimination, and increased negative self-perception and diminished social identity (Becares et al., 2009; Kirkbride et al., 2007; Pickett & Wilkinson, 2008; Shaw et al., 2012; Vogt Yuan, 2007). These are also relevant, to varying degrees, where there is a lack of fit between other examples of individual characteristics, such as family and socio-economic status, and the neighbourhood social environment. However other types of minority status have received relatively little attention. One study in Maastricht found the effect of being single on schizophrenia risk was almost doubled for people living alone in areas where living alone was less common than average (van Os *et al.*, 2000). Also a Swedish cohort study found that school children who were socio-economically deprived were more likely to experience psychotic symptoms in later life if they went to a school where others were on average less deprived (Zammit *et al.*, 2010). Ethnic density effects have also been shown for suicide and self-harm (Neeleman and Wessely, 1999, Neeleman *et al.*, 2001, Termorshuizen *et al.*, 2015) and also depression (Halpern and Nazroo, 2000, Propper *et al.*, 2005, Pickett *et al.*, 2009, Das-Munshi *et al.*, 2010). However, few recent studies have looked at these outcomes for other types of group density. There is, though, some evidence for an association between suicide and being in a minority due to socio-economic status, with higher rates shown for unemployed people living in low unemployment areas (Platt and Kreitman, 1985, Platt, 1986). Also a recent study looked at depression and being in a minority due to low socio-economic status but failed to find an association (Albor et al., 2014). A recent US survey also found lower rates of depression and anxiety among lesbian, gay and bisexual (LGB) respondents in states with a greater proportion of same sex couples (Hatzenbuehler et al., 2011).

Typically, though, studies of group density effects have analysed large datasets of individual health records linked to area level data. Relying on cases identified through psychiatric records alone risks diagnostic bias acting as a confounder. Where individual characteristics are at odds with their environment, it has been argued, they may be more likely to stand out and be picked up by mental health services (Wechsler and Pugh, 1967). Furthermore, given that the majority of mental disorders are untreated, it may be difficult to distinguish between patterns of help seeking behaviour and disease incidence, particularly for less severe disorders. Another problem with using routinely collected data is that it is usually not possible to determine length of residence, and distinguish between situations where people move to a particular type of neighbourhood as a result of mental ill health (‘social drift’) and those where the neighbourhood is itself a causal factor. Also, health records typically include very little information about the social characteristics of patients. One exception in recent years is ethnicity which is now routinely collected in UK health records, as it is in some other countries, and this may in part have contributed to the recent focus on ethnic density.

With this study we set out to test the hypothesis that different types of minority status are associated with increased risk of mental disorder. Our secondary aim was to compare the relative effects of different types of minority status on a range of outcomes in order to learn more about possible mechanisms behind these effects. In order to overcome the limitations of studies based on

health records alone, as outlined above, we used community survey data collected as part of the South East London Community Health (SELCoH) study. This is a large scale community psychiatric and physical morbidity survey that includes detailed information about the social characteristics of participants including length of residence (Hatch *et al.*, 2011). By linking these data with census data for the local neighbourhood we were able to examine the effect of minority status, defined by: ethnic group (Black African or Black Caribbean), household status (living alone) and occupational social class (disadvantaged) on a range of mental disorders. We were able to then relate this to three outcomes: psychotic experiences, common mental disorders and parasuicide.

## Methods

### Sample

We used data collected from the initial phase of SELCoH, comprising 1,698 individuals in 1,075 households randomly selected from the London boroughs of Lambeth and Southwark between 2008 and 2010. Both areas are ethnically diverse and include localities with widely differing socio-economic profiles. The overall sample was similar to the 2011 UK Census in terms of the demographic profile, particularly ethnicity and socioeconomic indicators, for this area (ONS, 2012). For a detailed comparison see (Morgan *et al.*, 2014).

### Outcomes

We assessed sub-clinical psychotic experiences using the Psychosis Screening Questionnaire (PSQ) (Bebbington and Nayani, 1995). The PSQ is structured as a set of initial probe questions followed by secondary questions. Following previous studies that looked at psychotic experience alone, we excluded question domains related to hypomania and defined psychotic experience as any positive response to secondary questions from the remaining domains (Morgan *et al.*, 2009, Morgan *et al.*, 2014). The presence of a common mental disorder, such as depression or anxiety based disorders, was defined as a score of 12 or more on the revised Clinical Interview Schedule (CIS-R) questionnaire (Lewis *et al.*, 1992). Parasuicide was determined by self-report, in response to the survey question: "Have you ever made an attempt to take your life, by taking an overdose of tablets or in some other way?"

### Predictors

Minority status was derived by comparing each individual social characteristic with the prevalence of the same characteristic in the local neighbourhood, defined as the nearest census lower super output area (LSOA). This is the most detailed area level at which relevant UK census data is available and corresponds to, on average, around 1500 people. Previous studies have shown ethnic density effects to be clearest at this detailed area level (Schofield *et al.*, 2010, 2016).

Ethnicity was based on self-reported identification with either Black Caribbean or Black African census defined ethnic groups. These are the two largest minority ethnic groups in this area, in Lambeth (11.6% Black African; 9.5% Black Caribbean) and Southwark (16.4% Black African; 6.2% Black Caribbean)(Office for National Statistics, 2012). However, the number of survey participants in these groups was still small (143 Caribbean and 234 African) therefore, to improve statistical power, we collapsed both groups, in the main analysis, to create an overall Black category. The equivalent area ethnic density measure was derived by adding the proportion of census derived Black Caribbean and Black African residents in each LSOA.

Single household status was based on the number of people living in each surveyed household and, at area level (LSOA), the proportion of people in single households was derived from recent census (2011) data.

Occupational social class was defined using the Registrar General's social class scale, based on current occupation (Office of Population Censuses and Surveys, 1980). This was dichotomised into advantaged (professional, managerial and technical, skilled non-manual and skilled manual) and disadvantaged (partly-skilled, unskilled and unemployed) groups. The following were excluded as they could not be easily assigned to either group: student, temporary sick/disabled, retired and looking after children. At area level (LSOA), occupational social class was defined using the National Statistics Socio-Economic Classification (NS-SEC) (Chandola and Jenkinson, 1999). This was split into two categories to match the individual level measure: advantaged (higher managerial, lower managerial and intermediate) and disadvantaged (lower supervisory, semi-routine, routine and never worked / unemployed). We excluded small employers and own account workers as these could not be easily classified.

We adjusted for area deprivation using the Index of Multiple Deprivation (IMD 2010). The IMD is derived from national data collected on seven domains (income, employment, education, health, crime, barriers to housing and services and living environment) (Department for Communities and Local Government, 2011). We also adjusted for age and gender in each analysis, with age entered in the following categories: 16-24, 25-34, 35-44, 45-54, 55-64, and 65+ years to account for non-linear effects.

## Ethical Approval

Ethical approval for the survey was granted by King's College London Research Ethics Committee (CREC/07/08-152)

## Statistical analysis

The main analysis was conducted using multilevel logistic regression to simultaneously account for effects at three levels: 1) individual respondent 2) household (to account for survey design) and 3) neighbourhood (LSOA). Minority status was operationalised, for each characteristic in turn, as a cross-level interaction between each individual characteristic and the area level prevalence of that characteristic. The effect of minority status on each mental health outcome, derived from the regression coefficients, was expressed in terms of a 10% decrease in the area prevalence of each characteristic for respondents with that characteristic. This follows the approach of previous ethnic density studies (Das-Munshi *et al.*, 2010, 2012) and allows comparison across different types of minority status. SELCoH includes non-response weights to account for age and gender discrepancies between the study sample and the overall population sampled. When incorporating weights in a multilevel model it is usually recommended to use re-scaling methods for weights at lower levels (Pfeffermann *et al.*, 1998). However, as age and gender are together highly related to the study outcomes we chose not to rescale weights, but instead entered these at the individual level only, assuming equal probability of selection at higher levels (Rabe-Hesketh and Skrondal, 2006). As a sensitivity analysis we reran the analysis both with and without weights to see whether this made any difference to our overall conclusions. We carried out a further sensitivity analysis testing for the effect of possible social drift by re-running the analysis excluding all those who had moved within the past two years. We also re-did the analysis using disaggregated ethnic groups to see if this would make a difference. We adjusted for area level deprivation where relevant and all results were adjusted for age and gender.

All analyses were conducted using Stata version 14 (StataCorp, 2015).

## Results

### Sample characteristics

Our study sample included 1,698 respondents in 1,067 households and 322 lower super output areas. Of these respondents, all but 8 completed the psychotic experiences (PSQ) questionnaire, all but 6 completed the common mental disorders (CIS-R) questionnaire, and all but 12 responded to the parasuicide question. Ethnicity was recorded for all but two and household status for all but 4 respondents. For occupational social class the above exclusions, outlined above, meant that 247 (14.6%) were removed. Area level (LSOA) data was matched for all study respondents. A substantial minority of respondents (26%) had experienced of some form of mental distress, according to the measures we looked at, with 19% having had a psychotic like experience in the past year, 23% classified as currently having a common mental disorder (a CIS-R score of 12 or over), and 8% having attempted suicide at some point in their lives (table 1).

### Evidence for cross level interactions

We found evidence for a cross-level interaction when we included group density effects for: ethnicity and psychosis symptoms (likelihood ratio test  $p < 0.001$ ) and para-suicide (lrtest  $p < 0.001$ ); single household status and psychosis (lrtest  $p = 0.01$ ); and some (albeit weak) evidence for social class and para-suicide (lrtest  $p = 0.06$ ). We also found a significant difference for ethnic density and depression (lrtest  $p = 0.02$ ) although the within group effect was non-significant.

Table 1 goes here

### Ethnic density and psychosis symptoms

Taking each outcome in turn (table 2), we found a clear relation between ethnic density and psychotic experiences. We found that being Black in an area with 10% fewer Black people was associated with a significant increase in the odds of reporting a psychotic experience, odds ratio (OR) 1.34 (95% CI 1.07 to 1.67). When we looked at each ethnicity sub-group we found this was apparent, although not statistically significant ( $p = 0.136$ ), for the Black Caribbean group only, with increased odds of psychotic experience in areas where there were fewer Black Caribbean people (OR 1.99 CI 0.81 to 4.89) (appendix 1 table 3). However, when we looked at the overall proportion of Black African and Black Caribbean people combined, the effect for Black Caribbeans was much clearer (OR 1.48 CI 1.02 to 2.17,  $p = 0.04$ ) (appendix 1 table 4).

### Other group density effects on psychosis symptoms

We found some, albeit weak, evidence of an association ( $p = 0.082$ ) when we looked at living alone in an area with 10% fewer people who live alone and psychotic experiences (OR 2.18 CI 0.91 to 5.26). Being in a more disadvantaged social class in an area where this was less usual (10% less) failed to make a statistically significant difference to the odds of psychotic experiences ( $p = 0.153$ ) and in fact showed a small inverse relation (OR 0.88 CI 0.73 to 1.05).

### Group density and other outcomes

We found no association between the above measures of minority status and odds of common mental disorder. When we looked at parasuicide we found a clear association with ethnic density (OR 1.84 CI 1.19 to 2.85). Again when we looked at ethnicity sub-groups this effect was only retained

for the Black Caribbean group, and again only when we looked at the overall Black population in each area (OR 2.32 CI 1.22 to 4.42). We found no relation between being in a minority due to single household status and risk of parasuicide. However, being in a disadvantaged social class where this was less common was associated with greater odds of parasuicide (OR 1.33 CI 1.03 to 1.71). Area level deprivation did not make any difference to the results for ethnic density and occupational social class and was therefore removed from the final model. For the analysis of single household status, area deprivation made a small difference and was therefore retained as a covariate.

Table 2 goes here

### Sensitivity analyses

We conducted a sensitivity analyses designed to control for social drift by restricting the sample to respondents who had been at the same address for two years or more, removing approximately 35% of the original sample. We found this made very little difference other than reducing the statistical significance of the estimates (see appendix 2 table 5). We also conducted a sensitivity analysis to see if our results were influenced by the method we used to incorporate survey weights. The unweighted analysis made almost no difference to our overall results (see appendix 2 table 6).

## Discussion

### Summary

We found that the experience of being in a minority due to ethnicity, single household status and occupational social class was associated with some, but not all, of the mental health outcomes we looked at. We found a strong ethnic density effect on both risk of psychotic experiences and parasuicide. We also found that living alone in an area where this was less usual was associated with increased risk of psychotic experience and that being in a disadvantaged social class in an area where this was less usual was associated with greater risk of parasuicide. We found no evidence that these group density effects were associated with common mental disorders.

### Strengths and limitations

The community survey data we used gave us a unique opportunity to compare the effects of different types of minority status on a range of mental disorders in the same study. We were able to avoid the diagnostic bias that can occur when using health records alone while the richness of the data allowed us to look at a variety of experiences of minority status. This was, though, cross-sectional data only and, while we could address issues with reverse causation where people have recently moved, we are unable to say much about the housing history of our sample beyond the previous two years. Therefore, we cannot rule out selection bias for mental disorders lasting more than two years or parasuicide prior to this. We would, though, expect 'social drift' to work in the opposite direction, so that people suffering mental distress would be more likely to move into areas with a higher proportion of people in these minority groups.

A further limitation is that, despite the survey being large (n=1698), we were restricted to fairly crude definitions of minority status to achieve statistical power. We were though able to investigate, to a limited extent, ethnic density effects within ethnic sub-groups. Our definition of occupational social class was, by necessity, crude as we had to match classes at individual and area levels measured on different scales. Also comparison with the social class measure is hindered by the amount of missing data, 247 (15%) missing responses, whereas this was negligible for the other measures. Our measure of single household status is also, by necessity, crude. For example, it could

miss those who effectively live alone but are in shared households, due to economic necessity, although when we compared gross income for individuals living alone and those in shared households we found no overall difference.

When comparing types of minority status it is also important to bear in mind that their prevalence varied widely and this may influence the strength of the reported effects. For example, the proportion of people with single household status was small (13%) compared to the proportion of Black people (22%). Also the extent to which each status varies at neighbourhood level also differs. When comparing outcomes, caution is needed as the measures used are on different temporal scales: the CIS-R question items relate to the recent past, i.e. the past month and past week, whereas the psychotic experiences (PSQ) questionnaire relates to the previous year, and parasuicide is assessed over a lifetime. Also the parasuicide question is potentially more open to bias, compared to the other outcome questions, as respondents may find this particularly uncomfortable to recall or disclose to others. It is, though, worth noting that a previous study, comparing these responses with national survey data, found these were almost double the national estimates (Aschan et al., 2013).

Also with any survey sample it is difficult to rule out selection effects. For example, we were surprised that area deprivation made relatively little difference to the results but this may simply reflect this particular sample. Lastly, our study is restricted to a specific urban area, south east London although this has distinct advantages as the diverse nature of the area and the richness of the data collected allowed us to look at a variety of social situations at a detailed geographical level. This is particularly important in an urban area, such as London, where localities with very different socioeconomic and ethnic profiles are often in close proximity to each other.

## Comparison with previous research

Overall our results confirm what has previously been suggested from similar studies looking at different types of minority status. However, our results showing a clear ethnic density effect associated with psychotic experiences measured using the PSQ differ from a previous national survey which failed to find an effect for the Black Caribbean group (Das-Munshi *et al.*, 2012). It is possible that this is because they looked at a more dispersed national sample, of lower ethnic density areas, whereas we were able to look at a specific urban area with a high concentration of this ethnic group. It is notable that studies of schizophrenia incidence using detailed local data from the UK have shown very similar results to ours (Boydell et al., 2001; Kirkbride et al., 2007, 2014; Schofield et al., 2010). For example, one recent study in East London found that a 1-SD increase in own-group ethnic density was associated with reduced psychosis incidence (RR 0.70; CI 0.48 to 0.99) among Black Africans (Kirkbride et al., 2014).

Looking at the association with ethnic density and parasuicide our results are in line with findings from previous work on suicide in South London. Neeleman found that the relative risk of Black and minority ethnic (BME) suicide declined (relative risk 0.67 CI 0.51 to 0.87) for each standard deviation increase in BME density (Neeleman and Wessely, 1999). Looking further afield, a Dutch study investigated all suicides in the four largest cities in the Netherlands and found lower risk among non-Western people (RR = 0.72, P = 0.004) in neighbourhoods with high (over 56%) versus low non-Western minority density (under 37%).

The absence of a corresponding ethnic density effect on risk of common mental disorders is perhaps not surprising given previous research. A recent review found that, out of four UK studies of Black African and Caribbean populations, only one showed a positive ethnic density effect (Shaw *et al.*, 2012a). Our previous study did show an inverse relation between ethnic density and depression



diagnosis for Black Africans, although for Black Caribbeans this effect appeared to work in the opposite direction (Schofield *et al.*, 2016). The detrimental effect of ethnic density on the health of Black Caribbean people has been previously reported in general population studies in the UK (Bécares *et al.*, 2012; Stafford *et al.*, 2011). However, the present study was based on primary care health records collected in South and East London and, as we argued, results may therefore have reflected health behaviour rather than underlying disorder.

For the effect of single household status our results are very similar to the previous Maastricht study where schizophrenia risk was almost doubled in areas with below average rates of single people (RR 10.3 95% CI 5.6 to 19.2) compared to areas with above average rates (RR 4.2 95% CI 1.9 to 9.3) (van Os *et al.*, 2000). While we looked at a sub-clinical measure of psychotic experiences, the fact that our results so clearly mirror previous research on schizophrenia is perhaps not surprising as a number of studies have shown the PSQ to be similarly related to risk factors for clinical psychosis (Morgan *et al.*, 2009, Das-Munshi *et al.*, 2012).

One study in recent years has examined the effect of being marginalised by occupational social class. Albor *et al.* looked at a large sample (4871) of mothers from the UK Millennium Cohort Study and examined the effect of socio-economic status, based on education and occupation, on depression or anxiety (Albor *et al.*, 2014). They found that people with high status in high status neighbourhoods had lower odds of depression or anxiety compared to those with high status in low status neighbourhoods. However, they failed to find the converse effect, as we did, instead they did not find a neighbourhood effect for those in the low status group. Also Zammit *et al.* (Zammit *et al.*, 2010) used a composite measure of deprivation and found the association with schizophrenia risk was reduced ( $P=0.06$ ) for those living in neighbourhoods where deprivation was more common. That our study failed to replicate this, in fact showing a small effect in the opposite direction, may simply reflect the fact that deprivation is conceptually distinct from social class. Our results showing a relation between minority status due to social class and parasuicide do, however, have a parallel with a previous study of unemployment and parasuicide. Rates of parasuicide among unemployed people in Edinburgh were shown to decline, in relative terms, when the prevailing unemployment rate increased (Platt and Kreitman, 1985, Platt, 1986). It is also worth noting that a number of studies have also shown a link between being in a minority due to socio-economic status and poor physical health and increased mortality (Yen and Kaplan, 1999, Winkleby *et al.*, 2006, Albor *et al.*, 2009).

## Interpretation of results

These results suggest that there is something about being different from others in the local neighbourhood that can be deleterious to mental health. They provide support for some of the hypothesised mechanisms behind the ethnic density effect; that this is not simply about being a numerical minority, but about racialised identities and the protection from discrimination that can result from living with other ethnic minorities (Bécares *et al.*, 2009). One concern was that by combining two distinct ethnic groups, Black Caribbean and Black African, we risked losing much of what makes these groups coherent. In fact, we found that each had a distinct mental health profile, with 31% of Black Caribbeans reporting psychotic experiences compared to 22% for Black Africans. Therefore, it is perhaps not surprising that the ethnic density effects differed between sub-groups. What was particularly interesting was that this became much clearer when we looked at specific groups in relation to the broader category of Black people living locally. This suggests that it may not be proximity to one's own ethnic group but proximity to others from a visible ethnic minority that is

the relevant factor. As others have argued, it seems it is not ethnicity itself that is relevant here but rather the social situation in which ethnicity is defined (Pickett and Wilkinson, 2008). This would therefore be a useful area of investigation for future research.

The strong ethnic density effect we found does not preclude other forms of minority status also being relevant, as we have shown. It is though unclear why the observed minority social class effect on parasuicide was not also seen when we looked at psychotic experiences. It is also unclear why there is no association between parasuicide and being in a minority due to single household status. However, the much smaller area variation in single household status (IQR 5% for single households compared to IQR 16% for percentage of Black people) may partly explain the absence of an effect for a relatively rare outcome such as parasuicide. It may also be that living alone is simply a more transitory state compared to occupational social class and therefore less likely to be related to a lifetime measure of parasuicide.

Furthermore, these examples of minority status are related. For example, those who are isolated in terms of ethnicity may also be isolated in terms of occupational class. We did, though, adjust for ethnicity in subsequent analyses (see appendix 3 table 7) and this made very little difference to the overall results. Also the outcomes may themselves be inter-related. However, when we adjusted for other outcomes in the analysis this made little difference to the reported results (appendix 3 table 8). It is also possible that those who have become unemployed as a result of mental ill health then find themselves in a minority in the area where they live and this could act as a confounder. We therefore also adjusted for the independent effect of being unemployed (appendix 3 table 9) but again found this made no difference to the overall results.

Looking at possible theoretical explanations for these differences; the experience of living alone, where this is less usual, may lead to a greater sense of being disconnected from others and therefore be more likely associated with psychotic experiences such as paranoia. The negative effects of being in a minority due to social class may be more likely to result in a negative self-image, due to lower perceived social position, leading to a greater tendency to parasuicide rather than psychosis.

## Conclusion and study implications

This study has been able to widen the scope of research on the relation between group density and mental distress for a range of mental disorders. Having shown a strong ethnic density effect on psychotic experiences and parasuicide we have also shown that these effects are not unique to ethnicity but can apply to other situations where people are in a minority. Further work is now needed to explore the mechanisms behind these increased risks. For example, is it a lack of social support resulting from minority status that is most important or an increased vulnerability to discrimination, as some studies have proposed? Alternatively, is it the perception of minority status / difference that is most relevant? We therefore recommend further research looking at a range of different types of minority status to better elucidate mechanisms behind the psychosocial pathways leading to mental disorders. It would also be useful to be able to investigate the timing and duration of relevant neighbourhood exposures using longitudinal designs to better elucidate possible causal mechanisms.

## Acknowledgements

The study author, Peter Schofield, was funded as part of a UK Medical Research Council fellowship (grant number MR/K021494/1). SLH and MH receive salary support from the National Institute for

Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. LB was supported by a Hallsworth Research Fellowship and a UK Economic and Social Research Council Future Research Leaders grant (grant number ES/K001582/1).

This paper represents independent research funded by the Medical Research Council (MRC), National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the MRC, NHS, the NIHR or the Department of Health.

We also wish to acknowledge the contribution of Professor Mel Bartley who gave advice on the social class measures used in this study.

## Declaration of Interest

None.

## Bibliography

Albor, C., Pickett, K., Wilkinson, R. G. & Ballas, D. (2009). Are poor mothers and their infants healthier in richer areas? The protective effect of area socio-economic density. *Journal of Epidemiology & Community Health* 63, 91-91.

Albor, C., Uphoff, E. P., Stafford, M., Ballas, D., Wilkinson, R. G. & Pickett, K. E. (2014). The effects of socioeconomic incongruity in the neighbourhood on social support, self-esteem and mental health in England. *Social Science and Medicine* 111, 1-9.

Aschan L, Goodwin L, Cross S, Moran P, Hotopf M, Hatch SL. (2013) Suicidal behaviours in South East London: Prevalence, risk factors and the role of socio-economic status. *Journal of Affective Disorder* 150, 441-449

Bebbington, P. & Nayani, T. (1995). The Psychosis Screening Questionnaire. *International Journal of Methods in Psychiatric Research* 5, 11-19.

Becares, L. & Das-Munshi, J. (2013). Ethnic density, health care seeking behaviour and expected discrimination from health services among ethnic minority people in England. *Health Place* 22, 48-55.

Bécares, L. Nazroo, J., Jackson, J., & Heuvelman, H. (2012). Ethnic density effects among Caribbean people in the US and England: a cross-national comparison. *Social Science & Medicine*, 75(12), 2107-2115.

Becares, L., Nazroo, J. & Stafford, M. (2009). The buffering effects of ethnic density on experienced racism and health. *Health & Place* 15, 700-708.

Boydell, J., van Os, J., McKenzie, K., Allardyce, J., Goel, R., McCreadie, R. G. & Murray, R. M. (2001). Incidence of schizophrenia in ethnic minorities in London: ecological study into interactions with environment. *British Medical Journal* 323, 1336-1338.

Chandola, T. & Jenkinson, C. (1999). Social class differences in morbidity using the new U.K. National Statistics Socio-Economic Classification. Do class differences in employment relations explain class differences in health? *Annals of the New York Academy of Sciences* 896, 313-5.

- Das-Munshi, J., Bécaries, L., Boydell, J. E., Dewey, M. E., Morgan, C., Stansfeld, S. A. & Prince, M. J. (2012). Ethnic density as a buffer for psychotic experiences: findings from a national survey (EMPIRIC). *The British Journal of Psychiatry* 4, 282-90
- Das-Munshi, J., Becares, L., Dewey, M. E., Stansfeld, S. A. & Prince, M. J. (2010). Understanding the effect of ethnic density on mental health: multi-level investigation of survey data from England. *British Medical Journal* 341.
- Department for Communities and Local Government (2011). *English Indices of Deprivation 2010*. London.
- Halpern, D. & Nazroo, J. (2000). The Ethnic Density Effect: Results from a National Community Survey of England and Wales. *International Journal of Social Psychiatry* 46, 34-46.
- Hatch, S. L., Frissa, S., Verdecchia, M., Stewart, R., Fear, N. T., Reichenberg, A., Morgan, C., Kankulu, B., Clark, J., Gazard, B., Medcalf, R., team, S. E. s. & Hotopf, M. (2011). Identifying socio-demographic and socioeconomic determinants of health inequalities in a diverse London community: the South East London Community Health (SELCoH) study. *BMC Public Health* 11, 861.
- Hatzenbuehler, M. L., Keyes, K. M., & McLaughlin, K. A. (2011). The protective effects of social/contextual factors on psychiatric morbidity in LGB populations. *International Journal of Epidemiology*, 40(4), 1071–1080.
- Kirkbride, J., Morgan, C., Fearon, P., Dazzan, P., Murray, R. & Jones, P. B. (2007). Neighbourhood-level effects on psychoses: re-examining the role of context. *Psychological Medicine* 37, 1413-25.
- Kirkbride, J. B., Jones, P. B., Ullrich, S., & Coid, J. W. (2014). Social deprivation, inequality, and the neighborhood-level incidence of psychotic syndromes in East London. *Schizophrenia Bulletin*, 40(1), 169–180.
- Lewis, G., Pelosi, A. J., Araya, R. & Dunn, G. (1992). Measuring psychiatric disorder in the community: a standardized assessment for use by lay interviewers. *Psychological Medicine* 22, 465-86.
- Morgan, C., Fisher, H., Hutchinson, G., Kirkbride, J., Craig, T. K., Morgan, K., Dazzan, P., Boydell, J., Doody, G. A., Jones, P. B., Murray, R. M., Leff, J. & Fearon, P. (2009). Ethnicity, social disadvantage and psychotic-like experiences in a healthy population based sample. *Acta Psychiatrica Scandinavica* 119, 226-35.
- Morgan, C., Reininghaus, U., Reichenberg, A., Frissa, S., Hotopf, M., Hatch, S. L. & Team, S. S. (2014). Adversity, cannabis use and psychotic experiences: evidence of cumulative and synergistic effects. *British Journal of Psychiatry* 204, 346-353.
- Neeleman, J. & Wessely, S. (1999). Ethnic minority suicide: a small area geographical study in south London. *Psychological Medicine* 29, 429-436.
- Neeleman, J., Wilson-Jones, C. & Wessely, S. (2001). Ethnic density and deliberate self harm; a small area study in south east London. *Journal of Epidemiology and Community Health* 55, 85-90.
- Office for National Statistics (2012). *2011 Census: Ethnic group, local authorities in England and Wales*.
- Office of Population Censuses and Surveys (1980). *Classification of occupations*. 1980 London.

- Pfeffermann, D., Skinner, C. J., Holmes, D. J., Goldstein, H. & Rasbash, J. (1998). Weighting for unequal selection probabilities in multilevel models. *Journal of the Royal Statistical Society Series B-Statistical Methodology* 60, 23-40.
- Pickett, K. & Wilkinson, R. (2008). People like us: ethnic group density effects on health. *Ethnicity and Health* 13, 321-334.
- Pickett, K. E., Shaw, R. J., Atkin, K., Kiernan, K. E. & Wilkinson, R. G. (2009). Ethnic density effects on maternal and infant health in the Millennium Cohort Study. *Social Science & Medicine* 69, 1476-83.
- Platt, S. (1986). Parasuicide and unemployment. *British Journal of Psychiatry* 149, 401-5.
- Platt, S. & Kreitman, N. (1985). Parasuicide and unemployment among men in Edinburgh 1968-82. *Psychological Medicine* 15, 113-23.
- Propper, C., Jones, K., Bolster, A., Burgess, S., Johnston, R. & Sarker, R. (2005). Local neighbourhood and mental health: evidence from the UK. *Social Science & Medicine* 61, 2065-83.
- Rabe-Hesketh, S. & Skrondal, A. (2006). Multilevel modelling of complex survey data. *Journal of the Royal Statistical Society Series a-Statistics in Society* 169, 805-827.
- Schofield, P., Ashworth, M. & Jones, R. (2011). Ethnic isolation and psychosis: re-examining the ethnic density effect. *Psychological Medicine* 41, 1263-9.
- Schofield, P., Das-Munshi, J., Mathur, R., Congdon, P. & Hull, S. (2016). Does depression diagnosis and antidepressant prescribing vary by location? Analysis of ethnic density associations using a large primary-care dataset. *Psychological Medicine*, 46, 1-9.
- Shaw, R. J., Atkin, K., Becares, L., Albor, C. B., Stafford, M., Kiernan, K. E., Nazroo, J. Y., Wilkinson, R. G. & Pickett, K. E. (2012a). Impact of ethnic density on adult mental disorders: narrative review. *British Journal of Psychiatry* 201, 11-19.
- Stafford, M., Bécares, L., & Nazroo, J. (2009). Objective and perceived ethnic density and health: findings from a UK general population survey. *American Journal of Epidemiology*, 170(4), 484-493
- StataCorp (2015). *Stata statistical software: Release 14*. StataCorp LP: College Station, TX
- Termorshuizen, F., Braam, A. W. & van Ameijden, E. J. C. (2015). Neighborhood ethnic density and suicide risk among different migrant groups in the four big cities in the Netherlands. *Social Psychiatry and Psychiatric Epidemiology* 50, 951-962.
- van Os, J., Driessen, G., Gunther, N. & Delespaul, P. (2000). Neighbourhood variation in incidence of schizophrenia. Evidence for person-environment interaction. *British Journal of Psychiatry* 176, 243-8.
- Veling, W., Susser, E., van Os, J., Mackenbach, J. P., Selten, J.-P. & Hoek, H. W. (2008). Ethnic Density of Neighborhoods and Incidence of Psychotic Disorders Among Immigrants. *American Journal of Psychiatry* 165, 66-73.
- Wechsler, H. & Pugh, T. F. (1967). Fit of individual and community characteristics and rates of psychiatric hospitalization. *American Journal of Sociology* 73, 331-338.
- Winkleby, M., Cubbin, C. & Ahn, D. (2006). Effect of cross-level interaction between individual and neighborhood socioeconomic status on adult mortality rates. *American Journal of Public Health* 96, 2145-2153.

Yen, I. H. & Kaplan, G. A. (1999). Neighborhood social environment and risk of death: Multilevel evidence from the Alameda County Study. *American Journal of Epidemiology* 149, 898-907.

Yuan, A. S. V. (2008). Racial composition of neighborhood and emotional well-being. *Sociological Spectrum* 28, 105-129.

Zammit, S., Lewis, G., Rasbash, J., Dalman, C., Gustafsson, J. E. & Allebeck, P. (2010). Individuals, schools, and neighborhood: a multilevel longitudinal study of variation in incidence of psychotic disorders. *Archives of General Psychiatry* 67, 914-22.

**Table 1 Description of study respondents**

	Ethnicity		Household status		Occupational class		<i>Overall</i>
	Black	Not Black	Single household	Larger household	Dis-advantaged	Not dis-advantaged	
Age - mean (standard deviation)	38 (16)	41 (17)	53 (18)	38 (16)	40 (16)	40 (15)	40 (17)
Gender - male %	40	44	40	44	38	47	44
Psychotic experiences - PSQ case %	26	17	24	18	24	16	19
Depression - CIS-R 12+ %	23	24	28	23	29	21	23
Parasuicide -%	6	8	14	7	12	5	8
Missing - frequency (%)		2 (0.1)		4 (0.2)		247 (15)	-
Neighbourhood Density - mean, inter-quartile range) <sup>a</sup>	22 (16)	-	18 (5)	-	32 (22)	-	
Total – frequency (%)	377 (22)	1319 (78)	212 (13)	1482 (87)	421 (29)	1030 (71)	1698 (100%)

<sup>a</sup>Percentage of people with this status in a neighbourhood

**Table 2. Minority status and mental distress**

	Psychotic experiences (PSQ)		Depression (CIS-R 12+)		Parasuicide	
	OR (95% confidence interval)	P value	OR (95% confidence interval)	P value	OR (95% confidence interval)	P value
Ethnic density (if Black then effect of 10% decrease in area proportion of Black people) <sup>a</sup>	1.34 (1.07 to 1.67)	0.010	1.13 (0.88 to 1.44)	0.327	1.84 (1.19 to 2.85)	0.006
Household status (if in single household then effect of 10% decrease in area proportion of people in single households) <sup>b</sup>	2.18 (0.91 to 5.26)	0.082	0.86 (0.36 to 2.03)	0.723	1.65 (0.49 to 5.53)	0.417
Social class (if disadvantaged then effect of 10% decrease in area proportion of disadvantaged) <sup>a</sup>	0.88 (0.73 to 1.05)	0.153	1.08 (0.88 to 1.33)	0.452	1.33 (1.03 to 1.71)	0.028

<sup>a</sup>adjusted for age and gender <sup>b</sup>adjusted for age, gender and area deprivation