



The Fort Collins Commuter Study

DOI:
[10.1111/ina.12533](https://doi.org/10.1111/ina.12533)

Document Version
Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):
Koehler, K., Good, N., Wilson, A., Mölter, A., Moore, B. F., Carpenter, T., Peel, J. L., & Volckens, J. (2018). The Fort Collins Commuter Study: Variability in Personal Exposure to Air Pollutants by Microenvironment. *Indoor Air*. <https://doi.org/10.1111/ina.12533>

Published in:
Indoor Air

Citing this paper
Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights
Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy
If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



DR. KIRSTEN KOEHLER (Orcid ID : 0000-0002-0516-6945)

DR. ANDER WILSON (Orcid ID : 0000-0003-4774-3883)

DR. JOHN VOLCKENS (Orcid ID : 0000-0002-7563-9525)

Article type : Original Article

Corresponding author mail id: John.Volckens@colostate.edu

The Fort Collins Commuter Study: Variability in Personal Exposure to Air Pollutants by
Microenvironment

Kirsten Koehler¹, Nicholas Good², Ander Wilson³, Anna Mölter^{2*}, Brianna F. Moore², Taylor
Carpenter², Jennifer L. Peel², John Volckens^{2,4}

1. Department of Environmental Health Sciences, Johns Hopkins University, 615 North Wolfe Street, Baltimore, MD, USA 21205
2. Department of Environmental and Radiological Health Sciences, Colorado State University, 1681 Campus Delivery, Fort Collins, CO, USA 80523
3. Department of Statistics, Colorado State University, 1877 Campus Delivery, Fort Collins, CO, USA 80523
4. Department of Mechanical Engineering, Colorado State University, 1374 Campus Delivery, Fort Collins, CO, USA 80523

* Present address: School of Environment, Education, and Development. University of Manchester, Manchester, M139PL UK

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/ina.12533

This article is protected by copyright. All rights reserved.

Abstract

This study investigated the role of microenvironment on personal exposures to black carbon (BC), fine particulate mass (PM_{2.5}), carbon monoxide (CO), and particle number concentration (PNC) among adult residents of Fort Collins, Colorado, U.S.A. Forty-four participants carried a backpack containing personal monitoring instruments for eight nonconsecutive 24-hour periods. Exposures were apportioned into five microenvironments: Home, Work, Transit, Eateries, and Other. Personal exposures exhibited wide heterogeneity that was dominated by within-person variability (both day-to-day and between microenvironment variability). Linear mixed-effects models were used to compare mean personal exposures in each microenvironments, while accounting for possible within-person correlation. Mean personal exposures during Transit and at Eateries tended to be higher than exposures at Home, where participants spent the majority of their time. Compared to Home, mean exposures to BC in Transit were, on average, 129% [95% confidence interval: 101% 162%] higher and exposures to PNC were 180% [101% 289%] higher in Eateries.

Key words: Personal exposure, microenvironment, air pollution, ultrafine particles, indoor air, traffic-related air pollution

Practical Implications: Americans spend 60-90% of their time indoors and within various microenvironments (e.g., home, work, eateries, etc.), yet data is lacking about how exposure to air pollution varies between such indoor microenvironments and when in transit. Personal exposure to multiple air pollutants varied substantially between these microenvironments and from day to day. Exposures in Transit and Eateries tended to be among the highest observed

and contributed disproportionately to integrated daily exposures compared to the amount of time spent in these microenvironments.

1. Introduction

Air pollution is a major contributor to the global burden of disease.¹ In the United States, outdoor air pollution is regulated and monitored through a network of fixed sites. Human exposure to air pollution, however, occurs primarily indoors.² Air pollution exposures within indoor microenvironments are derived from a combination of indoor sources (e.g. cooking, cleaning, movement of people) and infiltration of outdoor (ambient) air pollution indoors.³⁻⁵ The United States Environmental Protection Agency (US EPA) and World Health Organization acknowledge that short- and long-term health effects may be associated with exposure to poor quality indoor air,^{6,7} yet indoor air quality monitoring is rare due to the lack of regulations for most indoor microenvironments.

Many studies have provided evidence of an association between exposure to ambient air pollution, which is measured at fixed monitoring sites, and a wide variety of health effects including premature mortality,⁸ heart failure,⁹ stroke,¹⁰ diabetes mellitus,¹¹ and many more. However, generally weak correlations between outdoor air pollutant concentrations measured at fixed sites and personal exposures have been shown on timescales from days to a year¹²⁻¹⁶ and little is known about the role of exposures in different microenvironments.

Indoor air pollution concentrations often exceed ambient concentrations due to additional indoor sources and the relatively low levels of air exchange (dilution) indoors.^{3,17-²⁰} Many personal exposure studies, however, have been limited to a single microenvironment, due to limitations in monitoring technology.²¹ Single-microenvironment studies have

evaluated air pollutant concentrations at home,²²⁻²⁴ at office buildings,^{25,26} near roadways,^{27,28} or in transit.²⁹⁻³¹ Recent technological advances have enabled a more holistic view of individual exposures as they occur across microenvironments.^{19,32-36} Such studies may prove useful for the design of interventions that seek to reduce or mitigate pollutant exposures by targeting specific microenvironments, activities, or pollutants for exposure reduction.³⁷

The goals of this study were to evaluate the magnitude of personal exposures to fine particulate matter (PM_{2.5}), black carbon (BC), carbon monoxide (CO), and ultrafine particles (quantified by the metric particle number concentration [PNC]), as a function of microenvironment, in a panel of 44 adult participants in Fort Collins, Colorado, U.S.A. We compared personal exposures within and across microenvironments. Variance components analysis was used to examine the between- and within-person variability in microenvironmental exposures.

2. Methods

2.1. Personal Exposure Measurements

The study area, Fort Collins, Colorado is a mid-sized U.S. city with typical ambient PM_{2.5} levels at or below current, health-based regulatory standards established by the US EPA.³⁸ The city covers approximately 120 km² and has approximately 150,000 residents. Personal exposure data were collected as part of the Fort Collins Commuter Study.³⁹ Briefly, 45 non-smoking participants were recruited to carry backpacks containing lightweight, direct-reading exposure monitors for PM_{2.5} (pDR-1200, Thermo Scientific, Franklin, MA), BC (AE51, AethLabs, San Francisco, CA), and CO (T15n, Langan, Inc., San Francisco, CA). The backpack also contained a global positioning system (GPS) receiver (BT-Q1000XT, QStarz, Taipei, Taiwan) and a temperature, humidity and light sensor (MSR Electronics,

GmbH, Switzerland), and an accelerometer used to assess compliance. On a subset of monitoring days, the backpacks also contained an instrument to measure PNC (DiscMini, Testo, Lenzkirch, Germany); the subset was selected based on instrument availability and to maximize within-person replicates. For each sampling day, participants picked up the backpack from the study office in the evening and kept it with them while they completed a full day of their typical activities (including commuting to and from work) and then returned the backpack to the study office on the following morning. Participants were recruited to commute from their home to their workplace on 8 weekdays. On four days participants commuted by car (two days on a high traffic route and two days on a lower traffic route) and four days participants commuted by bicycle (two days on a high traffic route and two days on a lower traffic route) while carrying the backpack.³⁹ All sensors recorded data at intervals of 10-seconds or less. Each participant was asked to complete eight sampling days; some participants completed additional sampling days in the case of instrument failure (the range of sampling days per participant was between 1 and 11). Inclusion criteria were the following: 18 to 65 years old, valid driver's license, current non-smoker, commute at least 1.5 miles (2.4 km) from their home to their workplace and have no regular exposure to occupational dust and fumes (e.g., construction, manufacturing, agriculture). The study was conducted in all seasons; however, participants were not scheduled during official State of Colorado and U.S. Federal holidays. The Colorado State University Institutional Review Board approved all study procedures and participants provided written informed consent.

The protocol for personal, spatiotemporal exposure monitoring was similar to previous work,^{34,40} and details on the method can be found in Good et al.³⁹ Collected exposure data and GPS coordinates at 10-second resolution were downloaded into a geographic information system (Igor Pro, version 6.37) and processed with geocoded home

and work addresses to define a personal exposure track,³⁴ from which exposures were extracted and categorized for each participant. The microenvironments were manually assigned based on participant's location, a time-activity diary, and environmental metadata (temperature, light intensity, speed, heart rate, and motion). A custom graphical user interface was built to facilitate the microenvironment assignment. The interface mapped each participant's study day alongside their metadata time-series (e.g., abrupt changes in these values can help to identify when a participant moves from indoors to outdoors). Microenvironments were assigned using the interface by selecting measurements within areas of the map or measurements on the time-series and assigning the corresponding microenvironment using the time-activity diary for reference. For this analysis, exposure data between 9 pm to 9 pm local time were analyzed across five microenvironments that constituted the majority of location/activities identified: Home, Work, Transit (including any bicycling or driving that was recreational or for transportation), Eateries (including restaurants and coffee shops), and Other. The Other category includes any indoor locations that were not Home, Work or Eateries and when the participant was outdoors, but not in Transit, such as going for a walk or spending time in their yard. For this analysis, any walking during non-commute (i.e. between work and home) times was considered to be recreational and included in the Other category.

Sampling days were excluded for participants with missing GPS data (n=10 sampling days across 7 participants) because we could not determine microenvironments and for one participant where only a single sampling day was completed. Additionally, within a sampling day, if an instrument failure resulted in less than 22 of 24 hours of data collection, the values from that specific instrument were excluded. The resulting dataset contained 373 days of personal monitoring for 44 participants.

The direct-reading PM_{2.5} measurements were corrected with a gravimetric correction factor (i.e. the ratio of the gravimetric concentration over the time-weighted average measurements from the sensor over the same time period) determined from a filter measurement collected downstream of the sensor. The gravimetric limit of detection (LOD) was 31 µg; any filter mass measurements not exceeding the limit of detection were replaced by LOD/√2 in the calculation of the gravimetric mass concentration. The CO sensor concentrations were adjusted for temperature using the manufacturer's recommended calibration. The limits of detection for the PM, BC, and CO sensors were 1 µg m⁻³, 0.01 µg m⁻³, and 0.01 ppm, respectively. Any 10-second measures below the LOD were replaced by LOD/√2. We also investigated other imputation methods for values below the LOD without meaningful changes in our results (see Supplemental Information for details). None of the PNC measures were below the instrument limit of detection.

The Colorado Department of Public Health and the Environment maintains sites that routinely monitor ambient PM_{2.5} and CO with hourly resolution. Ambient air quality data were downloaded from the US EPA Air Quality System Data Mart from one location that monitored PM_{2.5} and another that monitored CO in Fort Collins, CO. Using the hourly ambient concentration data, 24-hour average ambient concentration from 9 pm to 9 pm on sampling days were calculated to match the personal sampling. The limit of detection for the ambient CO monitor was 0.5 ppm and all hourly average values below the LOD were replaced by 0.25. Ambient BC and PNC are not routinely monitored in Fort Collins.

2.2. Statistical Analysis

Descriptive statistics were assessed for exposure measures and for participants age and sex. Mean concentrations, over the time spent in each microenvironment and for the 24-hour period, across all study days were calculated for each participant. Violin plots with overlain boxplots of time-weighted average exposures in each of the microenvironments and for the 24-hour mean were created from all measures (all sampling days across all participants).

Daily integrated exposures were calculated by integrating concentrations over the time spent in the corresponding microenvironment or over the 24-hour sampling duration for each pollutant⁴¹. Daily mass-time ratio, the proportion of the integrated exposure attributable to each microenvironment divided by the fraction of time spent in that microenvironment,³⁵ were also created for each participant:

$$\text{mass} - \text{time ratio} = \frac{\text{integrated concentration}_{\text{ME/total}}}{\text{time}_{\text{ME/total}}} \quad (1)$$

where ME/total indicates the fraction of integrated concentration or time within a microenvironment over the total integrated concentration or time. The mass-time ratio is a useful metric for examining whether an individual's exposure in a given microenvironment contributes disproportionately to their 24-hr average exposure. A mass-time ratio equal to 1 indicates that the fraction of an individual's integrated daily exposure is equal to the fraction of time spent in that microenvironment (i.e., exposure in that microenvironment is indicative of their 24-hr average); values larger than 1 indicate that the fraction of their integrated exposure exceeded the time spent in that microenvironment (i.e., exposure in this microenvironment exceeds the expected value based on their 24-hr average) and values less than 1 indicate that the fraction of their integrated exposure was less than the time spent in that microenvironment.

A natural log transformation was applied to all time-weighted average exposures in each microenvironment, the 24-hour average, and the ambient concentration 24-hour average, resulting in exposures that were approximately normally distributed. Pearson's R correlation coefficients were calculated to evaluate correlation between pollutants within the same microenvironment and between ambient and personal PM_{2.5} and CO concentrations for the 24-hour sampling day. Statistical analyses were conducted using Matlab (MathWorks, Inc., Natick, MA) and SAS v9.4 (SAS Institute, Cary, NC).

Linear mixed models were developed to estimate the difference in mean personal exposures by microenvironment. The model is:

$$Y_{ijk} = \mu + b_i + d_{j(i)} + \beta_k + e_{ijk} \quad (\text{Model 1}),$$

where Y_{ijk} is the log-transformed time-weighted average pollutant concentration (BC, PM_{2.5}, CO, or PNC) in a microenvironment k for participant i and sampling day j , μ is the *overall* mean exposure for that pollutant, and e_{ijk} is the unexplained error. To account for potential within-subject and within-day cluster, we include a participant-specific random intercept b_i and a day-specific random intercept nested within participant $d_{j(i)}$. An unstructured covariance structure was specified. The regression coefficient β_k represents a fixed-effect for microenvironment k , where the 'Home' microenvironment was considered the reference and omitted from the model. Due to the transformation of the exposure, results are presented as mean percent difference compared to Home and 95% confidence intervals for the percent difference are presented.

We estimated the relative contribution of within-person and between-person variability in within-microenvironment integrated exposure using a variance components analysis. Since participants do not spend the same amount of time in each microenvironment,

the within-microenvironment integrated exposures were used. A variance component analysis was conducted on log-transformed integrated exposure values. A linear mixed model was developed to examine the variance components:

$$Z_{ij} = \mu + b_i + e_{ij} \quad (\text{Model 2}).$$

In this model Z_{ij} is the log-transformed integrated pollutant exposure (BC, PM_{2.5} mass, CO, or PNC) in a given microenvironment for participant i and sampling day j ; Z_{ij} is modeled as normally distributed with overall mean μ , person-specific intercept b_i assumed to be normally distributed with zero mean and variance σ_{btw}^2 (the between-person variability), and error e_{ij} that is assumed to be normally distributed with zero mean and variance σ_{with}^2 (the within-person variability). The variance components in each microenvironment and for the 24-hour integrated exposures were modeled separately. The intra-class correlation coefficient (ICC) was calculated as:

$$ICC = \frac{\sigma_{btw}^2}{\sigma_{with}^2 + \sigma_{btw}^2} \quad (\text{Equation 1})$$

and takes values between zero and one, where smaller values indicate that the within-person variability is high (relative to between-person variability).⁴²

3. Results

Overall, exposures for 44 participants who completed 373 sampling days were included in the analysis. Demographic information on study participants and sample sizes are listed in Table 1. Participant age varied from 22 to 61 years with a mean age of 37. The sample size for BC and CO are 327 and 339 sampling days per pollutant, respectively. The sample size is lower for PM_{2.5} (287 sampling days) due to missing filter data (n=10) or malfunctioning sensor (n=63) and for PNC (123 sampling days) because fewer instruments were available for deployment. Descriptive statistics for personal exposures and time spent in each microenvironment (taken across all participants and replicates) and the 24-hour geometric mean and geometric standard deviation are shown in Table 2. Participants spent

approximately 13 hours of their time at Home, 6.6 hours at Work, and a smaller fraction of time in the remaining categories. The majority of participants (90%) spent less than 10% (2.4 hours) of their time in Transit.

An illustrative example of a single participant's daily exposure track is shown in Figure 1, where the color of the bars on the map indicates the pollutant (BC in black and PM_{2.5} in blue; CO and PNC not shown) and the height of the bar indicates the relative concentration. These concentrations are also shown as a time-series (Figure 1; top panel) where color shading represents each microenvironment encountered (Home in red; Work in green; Transit in gray; Eatery in blue; and Other in orange). As was typical, there was high variability in participant exposures on different transit routes and in different microenvironments. Exposures tended to be low overnight at Home and low when at Work. Large increases in BC exposure during Transit (gray shaded portions of the time series) are visible and accompanied by only modest increases in PM_{2.5}. The relative increase in BC compared to PM_{2.5} is more apparent on the map, particularly near some intersections, where BC concentrations increased dramatically, while PM_{2.5} concentrations increased only slightly. Conversely, while walking in the natural area (green shaded regions on map; orange shading on the time series), PM_{2.5} exposures were elevated (possibly due to pedestrian and bike traffic on unpaved surfaces) and BC levels were low.

Violin plots with overlain boxplots of the mean concentrations of BC, PM_{2.5}, CO, and PNC in each microenvironment and the 24-hour time-weighted average are shown in Figure 2 for all sampling days across all participants. The 24-hour time-weighted average concentrations tended to be less than: 3 $\mu\text{g m}^{-3}$ for BC, 15 $\mu\text{g m}^{-3}$ for PM_{2.5}, 1 ppm for CO, and 2×10^4 # cm^{-3} for PNC. For all pollutants, mean concentrations at Work were the lowest

among the five microenvironments. Mean concentrations in Eateries were highly variable, but represented many of the highest observed exposures (along with exposures in Transit). Within-person percent change in mean concentrations compared to the Home microenvironment and 95% confidence intervals are listed in Table 3. Compared to the Home microenvironment, BC mean concentrations were 129% (95% CI: 101-162%) higher in Transit, 93% (58-137%) higher in Eateries and 24% (8-44%) higher in Other microenvironments. PM_{2.5} mean concentrations were 70% (66-74%) lower at Work, 14% (1-25%) lower in Transit, and 39% (29-47%) lower in Other compared to Home, but not different while at Eateries (-13-33%). Mean concentrations of CO were elevated in Transit and lower at Work and Other compared to Home. PNC mean concentrations were elevated in Transit and Eateries and lower at Work compared to Home. On average, PNC mean concentrations were 180% larger (101-289%) in Eateries compared to the Home microenvironment (the largest within-person difference detected).

The average contributions of these microenvironments to mean daily PM_{2.5} exposures are shown by participant in Figure 3. The large day-to-day variability that existed for many participants is exemplified for Participant 42 in the inset, where the contribution of the different microenvironments is shown for each of the eight sampling days. The 24-hour average exposures for each sampling day, by participant, are shown in the right panel. The variation in daily PM_{2.5} mean concentrations was large both within and between participants. However, the within-person variability of integrated exposure tended to exceed the between-person variability. Intra-class correlation coefficients were between 0 and 0.49 for all pollutants and microenvironments (Table S1). ICC values were smallest for PNC for the 24-hour average. Large within- and between-participant variability was also observed for the other pollutants (see Supplementary Material Figures S1-S3). Participants with the highest

mean 24-hour $PM_{2.5}$ exposures did not necessarily have the highest exposures to the other pollutants. Pearson's R correlation coefficients were calculated for the 24-hour mean and microenvironment means between each pair of pollutants (see Supplementary Material Table S2; results were similar for Spearman's correlation coefficients). Correlations tended to be highest between $PM_{2.5}$ and BC and weakest between CO and PNC. The highest correlation observed was between $PM_{2.5}$ and BC at Eateries and weakest between CO and PNC at Eateries. Correlations at Home varied between 0.06 and 0.50.

The mass-time ratio represents the proportion of integrated exposure attributable to each microenvironment divided by the fraction of time spent in that microenvironment.³⁵ Violin plots, with overlain boxplots, of the mass-time ratio in each microenvironment are shown in Figure 4. At Home, mass-time ratios tended to be somewhat less than 1 for BC and somewhat greater than 1 for $PM_{2.5}$, CO, and PNC. Mass-time ratios at Work tended to be less than 1 for all pollutants. Although Transit typically accounted for less than 7% of participant's time per day (75th percentile of time), this microenvironment contributed disproportionately to BC, CO and PNC integrated exposure. Up to 71% of some individual's daily BC integrated exposure occurred in Transit (mass-time ratio up to 11). Eateries also tended to contribute a disproportionate fraction to integrated exposures for all four pollutants. Mass-time ratios for the Other microenvironments tended to center around 1, with higher variability compared to Home and Work.

We examined the association between ambient (outdoor) and personal exposure for both $PM_{2.5}$ and CO (both log-transformed) on a daily basis (24-hr averages) and found that the correlation (Pearson's R) was 0.30 and 0.40, respectively. Figure 5 shows the relationship between ambient concentrations and personal exposures (without log-

transformation). During the days of our study (9/12/2012 to 2/4/2014), daily average ambient measures of PM_{2.5} and CO were 7.1 $\mu\text{g m}^{-3}$ and 0.34 ppm, respectively. Personal exposures showed a wider range of concentrations for both pollutants than was observed in the ambient measurements.

4. Discussion

Few studies have examined personal exposure to multiple pollutants as a function of microenvironment due to challenges associated with monitoring; historically, personal sampling has been time-integrated across 8- or 24-hr durations (e.g. filter-based analysis of PM_{2.5} mass). Relatively little is known about the variability of personal exposure to air pollution within- and between- microenvironments. This study, with a median of nine 24-hr measurements per person, is one of the largest studies of personal, multi-pollutant exposure by microenvironment to date. The proportion of time spent by participants in each microenvironment was roughly consistent with national averages for working adults.⁴³ In comparison to The National Human Activity Pattern Survey (NHAPS), participants in this study typically (mean values reported) spent somewhat less time per day at Home (13.0 hours compared to 16.6 hours), a similar amount of time at Work as was reported for working adults (6.6 hours compared to 6.5 hours), a similar amount of time in Transit (1.2 hours compared to 1.6 hours), and less time in Eateries compared to time in bars/restaurants reported by those who visited that microenvironment (0.7 hours compared to 1.8 hours).²

Variance-components analysis (Table S1) suggests that within-person variability dominates integrated exposure in a given microenvironment (i.e., all ICC values were between 0 and 0.49); this variability is also evident in Figure 3 (inset) where the magnitude and relative contribution (to daily integrated exposure) of each microenvironmental exposure

can change considerably from one monitoring day to the next. Large within-person variability was also observed for the daily measures, as shown in the right-hand-side of Figure 3, where an individual's daily personal mean concentration often spans over a factor of five across repeated measurement days. Our results demonstrate lognormal variations in personal exposure between subjects, within subjects, and across the various microenvironments encountered. These results have implications for exposure modeling, because capturing such variability may prove challenging using traditional modeling approaches to capture short-term exposures (e.g., temporal land-use regression). These results also reinforce the need for repeated measures when seeking to describe an individual's personal exposure to air pollution; a single, 24-hr measurement can vary by as much as an order of magnitude from one day to another.

We are unaware of any other studies that have considered variance components across different microenvironments. Recent studies have examined the variance components for daily averaged PM_{2.5}, microbial agents, and volatile organic carbon.⁴⁴⁻⁴⁷ In a study in Gothenburg, Sweden,⁴⁵ daily PM_{2.5} concentrations measured among 29 subjects (less than 2 samples per participant, total n=43) variance was partitioned within- and between-participants and resulted in an ICC value equal to 0.62. ICC values tended to be less than 0.5, and approached zero for some elements, in PM_{2.5} trace analysis.⁴⁵ ICC values (with variance partitioned within- and between- participants) less than 0.5 were also observed for 24-hour PM_{2.5} exposures among adults in Hong Kong,⁴⁷ and for exposure to microbial agents in buildings.⁴⁴ Rappaport and Kupper used a nested design with person nested within city (no person is in both cities) to examine 24-hour VOC exposures in the United States and found that typically half or more of the variance was within-person.⁴⁶ A large proportion of the 24-hour average PM_{2.5} variability was also within-person among elderly non-smoking

participants in Helsinki, Finland (ICC=0.34) and Amsterdam, Netherlands (ICC=0.11).⁴⁸

High microenvironmental variability was also observed in a study of ultrafine particulate exposures among adult participants in Copenhagen, Denmark.⁴¹ The low values of ICC observed for Transit may suggest that even with eight replicates, between-day variability may still impact the ability to evaluate exposure-health relationships.⁴⁹ Even when our dataset was restricted to sampling days for which participants commuted to work by car, ICC values were somewhat smaller for BC, PM_{2.5} and CO than observed for the full dataset including bicycle commuting, but somewhat larger for PNC. ICC values in Transit and Eateries were less than 0.35; however, the times spent in Transit and Eatery microenvironments were generally small for our participants.

Exposure levels were generally low compared to the United States National Ambient Air Quality Standards and WHO guidelines for indoor air for PM_{2.5} and CO. Mean microenvironment exposure levels were below the 24-hr WHO guideline of 25 $\mu\text{g m}^{-3}$ for PM_{2.5} 91% of the time at Home, 98% of the time at Work, 97% of the time in Transit, and 94% of the time for the 24-hour average. Only one Work average exceeded the 8-hr WHO guideline for CO of 8.7 ppm and the 24-hr guideline of 6.1 ppm was not exceeded for any of the 24-hour averages. Air quality regulations for BC and PNC do not exist in the United States and the WHO has not established guidelines for these pollutants. Ambient air quality in Colorado is generally below regulatory limits, except for higher levels of ozone during summer months that occasionally exceed the 24-hour ozone standard.³⁸ During the days of our study (9/12/2012 to 2/4/2014), daily average ambient measures of PM_{2.5} and CO were 7.1 $\mu\text{g m}^{-3}$ and 0.34 ppm, respectively. Ambient BC and PNC are not routinely monitored in Fort Collins.

For many participants, the majority of their time is spent at Home, and thus their integrated exposures tend to be dominated by exposures in the Home (see Table 2 and Figure 3). Compared to integrated exposures, mass-time ratios can be helpful for determining which microenvironments contribute disproportionately to personal exposures to identify potential ways to reduce exposure. Large mass-time ratios for the Transit and Eatery microenvironments were expected due to the close proximity of participants to traffic and cooking sources, respectively. Larger mass-time ratios in Transit compared to Home were observed for BC and CO, but not for PM_{2.5} because BC and CO are considered stronger markers for traffic-related air pollution. These microenvironments with high mass-time ratios could be targets for future interventions studies seeking to reduce exposures. Rea et al.⁵⁰ found that the fractional contribution of Transit to integrated PM_{2.5} exposure was higher than the fraction of time spent in that microenvironment among elderly adults in Baltimore, Maryland and Fresno, California (i.e. mass-time ratio greater than one), but in this study mass-time ratios centered near one for PM_{2.5} in Transit. Branis and Kolomaznikova³⁵ reported personal PM_{2.5} exposures for a single participant in the Czech Republic over 239 days and calculated mass-time ratios. The highest concentrations were observed in restaurants and bars (where smoking was allowed) with an average mass-time ratio greater than 20. Typical mass-time ratios in the Eatery category were smaller for our study, likely due to indoor smoking bans in the United States, but values as high as 23 for PM_{2.5} and as high as 57 for PNC were observed. Mass-time ratios at Home were larger in this study than observed by Branis and Kolomaznikova (mean value in this study was 1.2 for PM_{2.5}, compared to 0.62), but smaller than those observed by Levy Zamora et al.³⁶ in a population of mostly non-working, women in Texas, U.S.A. The difference in the Home mass-time ratios may reflect the fact that participants in this study tended to work, but in relatively in clean environments

with mass-time ratios generally less than 1 at Work. Additionally, few of our participants reported any exposure to secondhand smoke.

There are a number of strengths to the study design. Each participant was monitored for three pollutants (PM_{2.5} mass, BC, and CO) and a subset of participants were additionally monitored for PNC exposure, on approximately 8 days resulting in one of the largest datasets of personal, spatiotemporal exposure for multiple pollutants. In this study, participants wore chest-mounted accelerometers and the backpack contained an accelerometer and a GPS unit, which allowed us to assess participant compliance with wearing the backpack during waking hours while participants were moving. Compliance was assessed by matching a threshold level of movement (estimated to represent a spatial change) by the participant (determined from a chest-mounted accelerometer) to the backpack accelerometer in two-minute windows. On average, 66% (standard deviation 22%) of participant movements were matched by backpack movement during each 24-hour measurement period. Compliance was highest in the Transit microenvironment (96%) and lowest in Home (53%) microenvironment. Participants were asked to carry the backpack containing monitoring instruments with them as they moved between rooms, but were not asked to carry the backpack while moving around a room. The potential impacts of participant non-compliance on our results is unclear.

Some limitations of the study include the representativeness of our study population and study design. Participants were non-smokers who lived within the city of Fort Collins and lived at least 1.5 miles from their workplace and were asked to begin their commute during peak commute hours (between 0700 – 0900 h in the morning and 1630-1800 h in the evening). Thus, our results may not be representative of individuals who telecommute, are unemployed or retired. Participant commutes were scripted (mode and route), splitting days

between driving and bicycling and driving on higher and lower trafficked routes. Our previous work showed that participants experienced higher exposures to PM_{2.5}, BC, and PNC while cycling compared to driving and that they may reduce their exposures to BC and CO by using an alternate commuting route.³⁹ This may have contributed to the within-person variability demonstrated in this microenvironment (see Table S1). Participants may have changed their behavior under this study design, and because they were required to carry a backpack, relative to their typical time activity patterns. Furthermore, only weekdays were sampled. The impacts on time spent in each microenvironment and on ICC are unclear. Participants were recruited from a wide range of occupations; however, none of our participants were employed in health care or food-preparation, and those with regular exposure to occupational dust or fumes (e.g., construction, manufacturing, agriculture) were not eligible for inclusion in the study. Additionally, this study may not be reflective of working adults as a whole, as the variability in exposures between urban, suburban, and rural environments is not well established. Furthermore, there is the potential for measurement error in the personal exposure measurements.

5. Conclusions

This study is among the largest to quantify time-resolved exposures to PM_{2.5}, BC, CO, and PNC and to apportion those exposures into five common microenvironments. Despite representing a small proportion of participant's time, mean exposures to PNC, BC, and CO in Transit were 32%, 129%, and 51% higher than Home exposures, respectively. Exposures, particularly to PNC, were higher in Eateries than other microenvironments and were highly variable. The relative importance of some microenvironments, like Transit and Eateries, to integrated exposures were apparent, with mass-time ratios as high as 20 observed in Transit and as high as 57 in Eateries. Variance component analyses showed that within-person, day-

to-day variability dominated the total variance in the dataset and exemplifies the importance of repeated measurements in epidemiologic studies of chronic personal exposure to air pollution.

Acknowledgements

The authors would like to thank Charis Ackerson, Maggie L. Clark, Kristen M. Fedak, Ashleigh Kayne, Christian L'Orange, Casey Quinn, Jessica Tryner, and Viney Ugave for their contributions to data collection and data processing. This work was funded by the United States Department of Health and Human Services (HHS), National Institute of Health (NIH), National Institute of Environmental Health Sciences (NIEHS) under grant ES020017 by the Center for Disease Control (CDC), National Institute for Occupational Safety and Health (NIOSH), Mountain and Plains Education and Research Center under grant OH009229. The content of this article is solely the authors' responsibility and does not necessarily represent the official views of the HHS, NIH, CDC, NIOSH, or MAP ERC. K. Koehler gratefully acknowledges support from the College of Veterinary Medicine and Biomedical Sciences College Research Council at Colorado State University. The authors have no conflict of interest to declare.

Table 1: Participant characteristics and sample sizes for personal monitoring of four pollutants.

<i>Participants (N=44)</i>	N (%) or mean (STD) or median (range)
Female	24 (55%)
Male	20 (45%)
Age (years)	37 (12)
<i>Personal Sampling Days</i>	
Total number of sampling days	373
Median number of observations per participant; range	9 (3-13)

Table 2: Mean and quartiles of time-weighted average exposure for BC ($\mu\text{g m}^{-3}$), PM_{2.5} ($\mu\text{g m}^{-3}$), CO (ppm), and PNC (particles/cm³) by microenvironment, among sampling days for which each microenvironment was visited, and for the 24-hour average.

	N	Geometric Mean	GSD	Min	25 th	50 th	75 th	Max
Home								
Hours per day	373	13.0	1.4	0.5	12.7	13.8	14.6	24.0
BC ($\mu\text{g m}^{-3}$)	326	0.34	2.25	0.02	0.19	0.35	0.60	3.21
PM _{2.5} ($\mu\text{g m}^{-3}$)	287	8.0	2.3	1.0	4.6	7.3	13.1	78.3
CO (ppm)	338	0.38	2.32	0.01	0.24	0.41	0.63	3.98
PNC ($\# \text{cm}^{-3} * 10^4$)	123	0.81	2.18	0.20	0.42	0.77	1.39	9.62
Work								
Hours per day	357	6.6	1.6	0.1	6.2	7.5	8.4	12.5
BC ($\mu\text{g m}^{-3}$)	313	0.28	2.48	0.01	0.17	0.29	0.50	16.40
PM _{2.5} ($\mu\text{g m}^{-3}$)	275	2.4	2.5	0.7	1.1	2.0	4.4	68.6
CO (ppm)	324	0.16	2.86	0.01	0.10	0.18	0.30	15.17
PNC ($\# \text{cm}^{-3} * 10^4$)	119	0.27	2.47	0.01	0.15	0.27	0.49	2.56
Transit								
Hours per day	373	1.2	1.7	0.2	0.9	1.3	1.7	4.1
BC ($\mu\text{g m}^{-3}$)	326	1.02	2.01	0.05	0.65	1.04	1.59	15.54
PM _{2.5} ($\mu\text{g m}^{-3}$)	287	6.8	2.2	0.7	4.3	6.9	10.8	95.0
CO (ppm)	338	0.58	2.18	0.01	0.38	0.58	0.85	10.52
PNC ($\# \text{cm}^{-3} * 10^4$)	122	1.07	2.02	0.18	0.66	1.02	1.50	22.75
Eatery								
Hours per day	108	0.7	3.1	0.04	0.3	0.8	1.6	7.1
BC ($\mu\text{g m}^{-3}$)	96	0.71	2.42	0.06	0.38	0.63	1.15	9.29
PM _{2.5} ($\mu\text{g m}^{-3}$)	90	8.4	3.5	0.7	4.5	8.4	19.9	130.3
CO (ppm)	95	0.41	2.81	0.01	0.24	0.41	0.79	2.94
PNC ($\# \text{cm}^{-3} * 10^4$)	36	2.26	4.26	0.13	1.03	1.64	5.35	53.45
Other								
Hours per day	285	1.5	3.1	0.04	0.8	1.8	3.0	17.0
BC ($\mu\text{g m}^{-3}$)	249	0.46	2.37	0.03	0.28	0.46	0.77	6.03
PM _{2.5} ($\mu\text{g m}^{-3}$)	219	4.9	2.9	0.7	2.5	5.5	8.4	245.0
CO (ppm)	259	0.30	3.14	0.01	0.19	0.32	0.61	5.77
PNC ($\# \text{cm}^{-3} * 10^4$)	86	0.70	2.76	0.06	0.36	0.63	1.30	19.38
24-Hour								
BC ($\mu\text{g m}^{-3}$)	327	0.43	1.94	0.03	0.29	0.42	0.65	3.54
PM _{2.5} ($\mu\text{g m}^{-3}$)	287	7.4	1.9	1.7	4.9	6.6	10.2	51.3
CO (ppm)	339	0.37	2.03	0.01	0.26	0.40	0.56	4.77
PNC ($\# \text{cm}^{-3} * 10^4$)	123	0.79	1.79	0.20	0.50	0.76	1.19	6.69

Table 3: Mean percent difference [95% confidence intervals for mean percent difference] in personal exposure in each microenvironment compared to the Home microenvironment.

	N	Home	Work	Transit	Eatery	Other
BC	1310	Ref.	-1 [-13, 14]	129 [101, 162]	93 [58, 137]	24 [8, 44]
PM_{2.5}	1158	Ref.	-70 [-74, -66]	-14 [-25, -1]	8 [-13, 33]	-39 [-47, -29]
CO	1354	Ref.	-57 [-62, -52]	51 [34, 70]	8 [-11, 30]	-22 [-32, -11]
PNC	468	Ref.	-67 [-74, -59]	32 [6, 64]	180 [101, 289]	-11 [-30, 13]

Figure 1: GPS trace for one sampling day for a single participant. Bar color denotes the pollutant (BC in black or PM_{2.5} in blue) and bar height denotes the relative concentration. The red star denotes the participant's home location and the green star denotes the participant's work location. The concentrations of BC and PM_{2.5} over time are shown in the top panel. The shading indicates the microenvironment in which the participant was located during that time period (Home in red; Work in green; Transit in gray; Eatery in blue; and Other in orange). Natural areas are denoted in green on the map.




Figure 2: Violin plots overlain with boxplots of time-weighted average concentration in each microenvironment and over 24-hours for BC, PM_{2.5}, CO, and PNC for all participants. The top and bottom line of each box shows the 25th and 75th percentile of exposures and the middle line shows the median. Whiskers on the boxplots show 95% coverage of the data; the axis is broken along the line to show outliers. Asterisks indicate that exposures in a given microenvironment were significantly different than in the Home microenvironment, when accounting for within subject autocorrelation.




Figure 3: Left: the contribution of the five microenvironments to mean exposures of PM_{2.5} mass (average for each participant; $\mu\text{g m}^{-3}$). Mean exposures to PM_{2.5} by microenvironment for all eight sampling days for a single participant are shown in the inset. Right: 24-hour average PM_{2.5} exposures for all sampling days by participant. Participants are ordered from lowest to highest in terms of integrated PM_{2.5} exposure.




Figure 4: Violin plots overlain with boxplots of mass-time exposure ratios in each microenvironment for BC, PM_{2.5}, CO, and PNC. The top and bottom line of each box shows the 25th and 75th percentile of exposures and the middle line shows the median. Whiskers on the boxplots show 95% coverage of the data; the axis is broken along the line to show outliers. The dashed line indicates a value of 1, meaning that the fraction of time in a microenvironment equals the fraction of daily integrated exposure in that microenvironment.





Figure 5: Scatter plot of mean 24-hour personal exposures and mean 24-hour ambient concentrations for PM_{2.5} and CO.



References

1. Cohen AJ, Brauer M, Burnett R, et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet*. 2017;389(10082):1907-1918.
2. Klepeis NE, Nelson WC, Ott WR, et al. The National Human Activity Pattern Survey (NHAPS): a resource for assessing exposure to environmental pollutants. *J Expo Anal Env Epid*. 2001;11(3):231-252.
3. Meng QY, Spector D, Colome S, Turpin B. Determinants of indoor and personal exposure to PM(2.5) of indoor and outdoor origin during the RIOPA study. *Atmos Environ*. 2009;43(36):5750-5758.
4. Meng QY, Williams R, Pinto JP. Determinants of the associations between ambient concentrations and personal exposures to ambient PM2.5, NO2, and O-3 during DEARS. *Atmos Environ*. 2012;63:109-116.
5. Abt E, Suh HH, Catalano P, Koutrakis P. Relative contribution of outdoor and indoor particle sources to indoor concentrations. *Environ Sci Technol*. 2000;34(17):3579-3587.
6. EPA. Introduction to Indoor Air Quality. 2018; <https://www.epa.gov/indoor-air-quality-iaq/introduction-indoor-air-quality>. Accessed July 12, 2018, 2018.
7. WHO. *WHO Guidelines for Indoor Air Quality: Selected Pollutants*. Geneva: World Health Organization; 2010.
8. Hoek G, Krishnan RM, Beelen R, et al. Long-term air pollution exposure and cardio-respiratory mortality: a review. *Environ Health-Glob*. 2013;12.

9. Shah ASV, Langrish JP, Nair H, et al. Global association of air pollution and heart failure: a systematic review and meta-analysis. *Lancet*. 2013;382(9897):1039-1048.
10. Shah ASV, Lee KK, McAllister DA, et al. Short term exposure to air pollution and stroke: systematic review and meta-analysis. *Bmj-Brit Med J*. 2015;350.
11. Janghorbani M, Momeni F, Mansourian M. Systematic review and metaanalysis of air pollution exposure and risk of diabetes. *Eur J Epidemiol*. 2014;29(4):231-242.
12. Avery CL, Mills KT, Williams R, et al. Estimating Error in Using Ambient PM2.5 Concentrations as Proxies for Personal Exposures: A Review. *Epidemiology*. 2010;21(2):215-223.
13. Gulliver J, Briggs DJ. Personal exposure to particulate air pollution in transport microenvironments. *Atmos Environ*. 2004;38(1):1-8.
14. Molter A, Lindley S, de Vocht F, et al. Performance of a microenvironmental model for estimating personal NO2 exposure in children. *Atmos Environ*. 2012;51:225-233.
15. Janssen NAH, Hoek G, Brunekreef B, Harssema H, Mensink I, Zuidhof A. Personal sampling of particles in adults: Relation among personal, indoor, and outdoor air concentrations. *Am J Epidemiol*. 1998;147(6):537-547.
16. Brown KW, Sarnat JA, Suh HH, Coull BA, Spengler JD, Koutrakis P. Ambient site, home outdoor and home indoor particulate concentrations as proxies of personal exposures. *J Environ Monitor*. 2008;10(9):1041-1051.
17. Abt E, Suh HH, Allen G, Koutrakis P. Characterization of indoor particle sources: A study conducted in the metropolitan Boston area. *Environ Health Perspect*. 2000;108(1):35-44.
18. Breyse PN, Buckley TJ, Williams D, et al. Indoor exposures to air pollutants and allergens in the homes of asthmatic children in inner-city Baltimore. *Environ Res*. 2005;98(2):167-176.
19. Buonanno G, Stabile L, Morawska L. Personal exposure to ultrafine particles: The influence of time-activity patterns. *Sci Total Environ*. 2014;468:903-907.
20. Lai HK, Kendall M, Ferrier H, et al. Personal exposures and microenvironment concentrations of PM2.5, VOC, NO2 and CO in Oxford, UK. *Atmos Environ*. 2004;38(37):6399-6410.
21. Dons E, Panis LI, Van Poppel M, et al. Impact of time-activity patterns on personal exposure to black carbon. *Atmos Environ*. 2011;45(21):3594-3602.
22. Meng QY, Turpin BJ, Lee JH, et al. How does infiltration behavior modify the composition of ambient PM2.5 in indoor spaces? An analysis of RIOPA data. *Environ Sci Technol*. 2007;41(21):7315-7321.
23. Baxter LK, Clougherty JE, Paciorek CJ, Wright RJ, Levy JI. Predicting residential indoor concentrations of nitrogen dioxide, fine particulate matter, and elemental carbon using questionnaire and geographic information system based data. *Atmos Environ*. 2007;41(31):6561-6571.
24. Karotki DG, Beko G, Clausen G, et al. Cardiovascular and lung function in relation to outdoor and indoor exposure to fine and ultrafine particulate matter in middle-aged subjects. *Environment International*. 2014;73:372-381.
25. Mosqueron L, Momas I, Le Moullec Y. Personal exposure of Paris office workers to nitrogen dioxide and fine particles. *Occup Environ Med*. 2002;59(8):550-555.
26. Horemans B, Worobiec A, Buczynska A, Van Meel K, Van Grieken R. Airborne particulate matter and BTEX in office environments. *J Environ Monitor*. 2008;10(7):867-876.
27. McAdam K, Steer P, Perrotta K. Using continuous sampling to examine the distribution of traffic related air pollution in proximity to a major road. *Atmos Environ*. 2011;45(12):2080-2086.

28. Fruin S, Westerdahl D, Sax T, Sioutas C, Fine PM. Measurements and predictors of on-road ultrafine particle concentrations and associated pollutants in Los Angeles. *Atmos Environ*. 2008;42(2):207-219.
29. Suarez L, Mesias S, Iglesias V, Silva C, Caceres DD, Ruiz-Rudolph P. Personal exposure to particulate matter in commuters using different transport modes (bus, bicycle, car and subway) in an assigned route in downtown Santiago, Chile. *Environ Sci-Proc Imp*. 2014;16(6):1309-1317.
30. Kam W, Delfino RJ, Schauer JJ, Sioutas C. A comparative assessment of PM_{2.5} exposures in light-rail, subway, freeway, and surface street environments in Los Angeles and estimated lung cancer risk. *Environ Sci-Proc Imp*. 2013;15(1):234-243.
31. Wang X, Gao HO. Exposure to fine particle mass and number concentrations in urban transportation environments of New York City. *Transport Res D-Tr E*. 2011;16(5):384-391.
32. Mazaheri M, Clifford S, Jayaratne R, et al. School Children's Personal Exposure to Ultrafine Particles in the Urban Environment. *Environ Sci Technol*. 2014;48(1):113-120.
33. Allen R, Wallace L, Larson T, Sheppard L, Liu LJS. Estimated hourly personal exposures to ambient and nonambient particulate matter among sensitive populations in Seattle, Washington. *J Air Waste Manage*. 2004;54(9):1197-1211.
34. Adams C, Riggs P, Volckens J. Development of a method for personal, spatiotemporal exposure assessment. *J Environ Monitor*. 2009;11(7):1331-1339.
35. Branis M, Kolomaznikova J. Year-long continuous personal exposure to PM_{2.5} recorded by a fast responding portable nephelometer. *Atmos Environ*. 2010;44(24):2865-2872.
36. Levy Zamora M, Pulczynski JC, Johnson N, et al. Maternal exposure to PM_{2.5} in south Texas, a pilot study. *Sci Total Environ*. 2018;628-629:1497-1507.
37. Egeghy PP, Quackenboss JJ, Sandra CC, Ryan PB. Determinants of temporal variability in NHEXAS-Maryland environmental concentrations, exposures, and biomarkers. *J Expo Anal Env Epid*. 2005;15(5):388-397.
38. Air Pollution Control Division. *2014 Air Quality Data Report*. Denver, CO: Colorado Department of Public Health and Environment,; March 2016 2016.
39. Good N, Molter A, Ackerson C, et al. The Fort Collins Commuter Study: Impact of route type and transport mode on personal exposure to multiple air pollutants. *J Expos Sci Environ Epidemiol*. 2016;26(4):397-404.
40. Adams C, Rabinovitch N, Strand M, Marquart K, Riggs P, Volckens J. New Insights into Personal Exposure to Particulate Matter Air Pollution: A Case Study of Children with Asthma in Denver. *Proc Am Thorac Soc*. 2010;7(2):152.
41. Beko G, Kjeldsen BU, Olsen Y, et al. Contribution of various microenvironments to the daily personal exposure to ultrafine particles: Personal monitoring coupled with GPS tracking. *Atmos Environ*. 2015;110:122-129.
42. Sahai H, Ageel MI. *The analysis of variance : fixed, random, and mixed models*. Boston: Birkhäuser; 2000.
43. Bureau of Labor Statistics. Table A-2. Time spent in detailed primary activities, and percent of the civilian population engaging in each detailed activity category, averages per day by sex on weekdays and weekends, 2013 annual averages. 2013. Accessed 10/20/14.
44. Cho SJ, Cox-Ganser JM, Kreiss K, Park JH. Evaluation of individual-based and group-based exposure estimation of microbial agents in health effects associated with a damp building. *J Expo Sci Env Epid*. 2013;23(4):409-415.

- Accepted Article
45. Johannesson S, Rappaport SM, Sallsten G. Variability of environmental exposure to fine particles, black smoke, and trace elements among a Swedish population. *J Expo Sci Env Epid.* 2011;21(5):506-514.
 46. Rappaport SM, Kupper LL. Variability of environmental exposures to volatile organic compounds. *J Expo Anal Env Epid.* 2004;14(1):92-107.
 47. Chen XC, Ward TJ, Cao JJ, et al. Determinants of personal exposure to fine particulate matter (PM_{2.5}) adult subjects in Hong Kong. *Sci Total Environ.* 2018;628-629:1165-1177.
 48. Lanki T, Ahokas A, Alm S, et al. Determinants of personal and indoor PM_{2.5} and absorbance among elderly subjects with coronary heart disease. *J Expo Sci Env Epid.* 2007;17(2):124-133.
 49. Brunekreef B, Noy D, Clausing P. Variability of Exposure Measurements in Environmental Epidemiology. *Am J Epidemiol.* 1987;125(5):892-898.
 50. Rea AW, Zufall MJ, Williams RW, Sheldon L, Howard-Reed C. The influence of human activity patterns on personal PM exposure: A comparative analysis of filter-based and continuous particle measurements. *J Air Waste Manage.* 2001;51(9):1271-1279.